

Datasheet: VMA00376KT

| Description: | ErbB4 ANTIBODY WITH CONTROL LYSATE | | |
|---------------|------------------------------------|--|--|
| Specificity: | ErbB4 | | |
| Format: | Purified | | |
| Product Type: | PrecisionAb™ Monoclonal | | |
| Clone: | 3A10 | | |
| Isotype: | lgG1 | | |
| Quantity: | 2 Westerns | | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol | | | | | |
|----------------------------|---|------------|----|----------------|--------------------|--|
| | recommendations, please visit www.bio-rad-antibodies.com/protocols. | | | | | |
| | Western Distline | Yes | No | Not Determined | Suggested Dilution | |
| | Western Blotting | - | | | 1/1000 | |
| | PrecisionAb antibodies have been extensively <u>validated for the western blot applic</u> antibody has been validated at the suggested dilution. Where this product has not been to use in a particular technique this does not necessarily exclude its use in such procedures optimization may be required dependant on sample type. | | | | | |
| Target Species | Human | | | | | |
| Product Form | Purified IgG - liquid | | | | | |
| Preparation | 20µl Mouse monoclonal antibody purified by affinity chromatography from ascites | | | | | |
| Buffer Solution | Phosphate buffered saline | | | | | |
| Preservative | 0.09% Sodium Azide (NaN | 3) | | | | |
| Stabilisers | 1% Bovine Serum Albumin | | | | | |
| | 50% Glycerol | | | | | |
| Immunogen | Recombinant protein fragment corresponding to aa 992-1308 of human ErbB4 (NP_005226) produced in <i>E.coli</i> | | | | | |
| External Database Links | UniProt: Q15303 Related rea | agents | | | | |
| | Entrez Gene: | | | | | |
| | | d reagents | | | | |
| Synonyms | HER4 | | | | | |

| Specificity | Mouse anti Human ErbB4 antibody recognizes the receptor tyrosine-protein kinase ErbB4, also known as ALS19, avian erythroblastic leukemia viral (v-erb-b2) oncogene homolog 4, proto-oncogene-like protein c-ErbB-4, receptor tyrosine-protein kinase erbB-4, HER4 and v-erb-a erythroblastic leukemia viral oncogene homolog 4. |
|----------------------------------|--|
| | The ERBB4 gene product is a member of the Tyr protein kinase family and the epidermal growth factor receptor subfamily. It encodes a single-pass type I membrane protein with multiple cysteine rich domains, a transmembrane domain, a tyrosine kinase domain, a phosphotidylinositol-3 kinase binding site and a PDZ domain binding motif. The protein binds to and is activated by neuregulins and other factors and induces a variety of cellular responses including mitogenesis and differentiation. Multiple proteolytic events allow for the release of a cytoplasmic fragment and an extracellular fragment. Mutations in ErbB4 have been associated with cancer. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized (provided by RefSeq, Jul 2008). |
| | Mouse anti Human ErbB4 antibody detects a band of 160 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. |
| Western Blotting | Anti ErbB4 detects a band of approximately 160 kDa in HEK293 cell lysates |
| Instructions For Use | Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page. |
| Lysate Composition | 400µg HEK293 lysate lyophilized in RIPA buffer |
| Lysate Reconstitution | If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. |
| Storage | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. |
| | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. |
| Shelf Life | As supplied, 12 months from date of despatch |
| Acknowledgements | PrecisionAb [™] is a trademark of Bio-Rad Laboratories |
| Health And Safety Information | Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u> |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M336942:181214'

Printed on 02 Jan 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint