

# Datasheet: VMA00381KT

Description:	CIP2A ANTIBODY WITH CONTROL LYSATE
Specificity:	CIP2A
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	OTI1C11
Isotype:	lgG1
Quantity:	2 Westerns

### **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Western Blotting	•			1/1000			
	PrecisionAb antibodies application. The antibodies product has not been test exclude its use in such p sample type.	dy has bee sted for us	en validato e in a par	ed at the suggested dil ticular technique this d	lution. Where this loes not necessarily			
Target Species	Human							
Product Form	Purified IgG - liquid							
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography from ascites							
Buffer Solution	Phosphate buffered salin	ne						
Preservative	0.09% Sodium Azide (N	aN <sub>3</sub> )						
Stabilisers	1% Bovine Serum Albur	0/						
	50% Glycerol							
Immunogen	Recombinant protein fra (NP_065941) produced	•	respondir	ng to aa 635-905 of hui	man CIP2A			
External Database	UniProt:							

Links	Q8TCG1 Related reagents
	Entrez Gene: <u>57650</u> KIAA1524 <u>Related reagents</u>
Synonyms	CIP2A
Specificity	<b>Mouse anti Human CIP2A antibody</b> recognizes CIP2A, also known as cancerous inhibitor of PP2A, cancerous inhibitor of protein phosphatase 2A and p90 autoantigen.
	Mouse anti Human CIP2A antibody detects a band of 90 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti CIP2A detects a band of approximately 90 kDa in HEK293 cell lysates
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg HEK293 lysate lyophilized in RIPA buffer
Lysate Reconstitution	<ul> <li>If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</li> <li>If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</li> <li>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</li> </ul>
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb <sup>™</sup> is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>
Regulatory	For research purposes only

## Related Products

#### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21	
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	From
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.march	

15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M350982:190315'

#### Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint