

Datasheet: VMA00393KT

Description:	RIP3 ANTIBODY WITH CONTROL LYSATE
Specificity:	RIP3
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Clone:	1D9
Isotype:	IgG2b
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography from ascites
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin 50% Glycerol
Immunogen	Recombinant protein fragment corresponding to aa 233-518 of human RIP3 (NP_006862) produced in <i>E.coli</i>
External Database Links	UniProt: Q9Y572 Related reagents Entrez Gene: 11035 RIPK3 Related reagents
Synonyms	RIP3

Specificity **Mouse anti Human RIP3 antibody** recognizes RIP3, also known as RIP-like protein kinase 3, receptor interacting protein 3 and receptor-interacting serine/threonine-protein kinase 3.

The product of the RIPK3 gene is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The RIP3 protein is predominantly localized to the cytoplasm, and can undergo nucleo-cytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor (provided by RefSeq, Jul 2008).

Mouse anti Human RIP3 antibody detects a band of 57 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Anti RIP3 detects a band of approximately 57 kDa in Jurkat cell lysates

Instructions For Use Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Lysate Composition 400µg Jurkat lysate lyophilized in RIPA buffer

Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.
- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.
Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.

Guarantee As supplied, 12 months from date of despatch

Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories

Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at:
Antibody (10048): <https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf>
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

'M338322:181217'

Printed on 15 Mar 2019
