

Datasheet: VMA00411KT

| Description: | CLATHRIN HEAVY CHAIN ANTIBODY WITH CONTROL LYSATE |
|---------------|---|
| Specificity: | CLATHRIN HEAVY CHAIN |
| Format: | Purified |
| Product Type: | PrecisionAb Monoclonal |
| Isotype: | lgG1 |
| Quantity: | 2 Westerns |
| | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | | | |
|-----------------------------|--|-------------------|-------------|--------------------------|------------------------------|--|--|--|
| | Western Blotting | Yes | No | Not Determined | Suggested Dilution 1/1000 | | | |
| | PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type. | | | | | | | |
| Target Species | Human | | | | | | | |
| Species Cross Reactivity | Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. | | | | | | | |
| Product Form | Purified IgG - liquid | | | | | | | |
| Preparation | 20µl Mouse monoclonal tissue culture supernatar | | ourified by | / affinity chromatograpl | ny on Protein G from | | | |
| Buffer Solution | Phosphate buffered salin | ie | | | | | | |
| Preservative Stabilisers | 0.09% Sodium Azide (Na | aN ₃) | | | | | | |

| Immunogen | <i>E.coli</i> -derived recombinant protein corresponding to aa 1521-1675 of human clathrin heavy chain | | | | | |
|----------------------------|---|--|--|--|--|--|
| External Database Links | UniProt: Q00610 Related reagents Entrez Gene: 1213 CLTC Related reagents | | | | | |
| Synonyms | CLH17, CLTCL2, KIAA0034 | | | | | |
| Specificity | Mouse anti Human clathrin heavy chain antibody recognizes clathrin heavy chain , also known as CLTC, CHC, clathrin heavy chain on chromosome 17 (CHC17) and CLTCL2. Clathrin is a major protein component of the cytoplasmic face of intracellular organelles, | | | | | |
| | called coated vesicles and coated pits. These specialized organelles are involved in the intracellular trafficking of receptors and endocytosis of a variety of macromolecules. The basic subunit of the clathrin coat is composed of three heavy chains and three light chains (provided by RefSeq, Jul 2008). | | | | | |
| | Mouse anti Human clathrin heavy chain antibody detects a band of 192 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. | | | | | |
| Western Blotting | Anti clathrin heavy chain detects a band of approximately 192 kDa in HepG2 cell lysates | | | | | |
| Instructions For Use | Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page. | | | | | |
| Lysate Composition | 400µg HepG2 lysate lyophilized in RIPA buffer | | | | | |
| Lysate Reconstitution | If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. | | | | | |
| Storage | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. | | | | | |
| | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. | | | | | |
| Guarantee | As supplied, 12 months from date of despatch | | | | | |
| Acknowledgements | PrecisionAb [™] is a trademark of Bio-Rad Laboratories | | | | | |

| Health A Informati | on Ant | Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u> | | | | | | | |
|-----------------------|------------------------|--|---|----------|--------------------------------------|--|--|--|--|
| Regulato | ry For | For research purposes only | | | | | | | |
| Relate | d Products | | | | | | | | |
| Recomm | nended Second | dary Antibodies | | | | | | | |
| Goat Ant | i Mouse IgG (H/L |) (STAR207) <u>HRP</u> | | | | | | | |
| Recomm | nended Negativ | ve Controls | | | | | | | |
| MOUSE Ig | G1 NEGATIVE CC | NTROL (MCA928) | | | | | | | |
| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 | | | | |
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 | | | | |
| | Email: antibody_sales_ | us@bio-rad.com | Email: antibody_sales_uk@bic | -rad.com | Email: antibody_sales_de@bio-rad.com | | | | |
| | | | no longer supply printed o access your digital versi 'M350981:190315' | | | | | | |
| | | F | Printed on 09 Feb 2021 | | | | | | |

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint