

Datasheet: VMA00417

Description:	MOUSE ANTI CORTACTIN
Specificity:	CORTACTIN
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG2a
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Human cortactin produced in HEK293 cells
External Database Links	<p>UniProt: Q14247 Related reagents</p> <p>Entrez Gene: 2017 CTTN Related reagents</p>
Synonyms	EMS1
Specificity	Mouse anti Human cortactin antibody recognizes the Src kinase substrate cortactin, also known

as CTTN, amplexin, ems1 sequence (mammary tumor and squamous cell carcinoma-associated (p80/85 src substrate) and oncogene EMS1.

The CTTN gene product is overexpressed in breast cancer and squamous cell carcinomas of the head and neck. The CTTN protein is localized in the cytoplasm and in areas of the cell-substratum contacts. CTTN has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of CTTN contributes to tumor cell invasion and metastasis. Three splice variants that encode different isoforms have been identified for CTTN (provided by RefSeq, May 2010).

Mouse anti Human cortactin antibody detects a band of 85 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti cortactin detects a band of approximately 85 kDa in MCF7 cell lysates
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Guarantee	12 months from date of despatch
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M370174:200529'

Printed on 11 Aug 2020