

Datasheet: VMA00417KT

Description:	CORTACTIN ANTIBODY WITH CONTROL LYSATE		
Specificity:	CORTACTIN		
Format:	Purified		
Product Type:	PrecisionAb Monoclonal		
Isotype:	lgG2a		
Quantity:	2 Westerns		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Western Blotting	-			1/1000			
	PrecisionAb antibodie <u>application.</u> The antibo product has not been te exclude its use in such p sample type.	dy has bee sted for us	en validate e in a par	ed at the suggested dil ticular technique this d	ution. Where this oes not necessarily			
Target Species	Human							
Product Form	Purified IgG - liquid							
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant							
Buffer Solution	Phosphate buffered saline							
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)							
Immunogen	Human cortactin produced in HEK293 cells							
External Database Links	UniProt: <u>Q14247</u> <u>Related</u>	reagents						

Entrez Gene:

2017 CTTN Related reagents

Synonyms	EMS1					
Specificity	Mouse anti Human cortactin antibody recognizes the Src kinase substrate cortactin, also known as CTTN, amplaxin, ems1 sequence (mammary tumor and squamous cell carcinoma-associated (p80/85 src substrate) and oncogene EMS1.					
	The CTTN gene product is overexpressed in breast cancer and squamous cell carcinomas of the head and neck. The CTTN protein is localized in the cytoplasm and in areas of the cell-substratum contacts. CTTN has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of CTTN contributes to tumor cell invasion and metastasis. Three splice variants that encode different isoforms have been identified for CTTN (provided by RefSeq, May 2010).					
	Mouse anti Human cortactin antibody detects a band of 85 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.					
Western Blotting	Anti cortactin detects a band of approximately 85 kDa in MCF7 cell lysates					
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.					
Lysate Composition	400µg MCF7 lysate lyophilized in RIPA buffer					
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H ₂ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT. - If using BME reconstitute the lyophilized lysate with 180 μ l DI H ₂ O, add 200 μ l 2x Laemmli					
	Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.					
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.					
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.					
Guarantee	As supplied, 12 months from date of despatch					
Acknowledgements	PrecisionAb [™] is a trademark of Bio-Rad Laboratories					
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>					

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...)<u>HRP</u> Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351004:190315'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint