

Datasheet: VMA00420KT

Description:	HIGH MOBILITY GROUP PROTEIN HMGI-C ANTIBODY WITH CONTROL LYSATE
Specificity:	HIGH MOBILITY GROUP PROTEIN HMGI-C
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

Target Species

Human

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

J .				
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.			
Product Form	Purified IgG - liquid			
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)			

Immunogen	E.coli-derived recombinant protein corresponding to aa 1-109 of human high mobility group protein HMGI-C				
External Database Links	UniProt: P52926 Related reagents				
	Entrez Gene: 8091 HMGA2 Related reagents				
Synonyms	HMGIC				
Specificity	Mouse anti Human high mobility group protein HMGI-C antibody recognizes high-mobility group protein HMGI-C also known as high mobility group AT-hook 2 and HMGA2.				
	The HMGA2 gene encodes a protein that belongs to the non-histone chromosomal high mobility group (HMG) protein family. HMG proteins function as architectural factors and are essential components of the enhanceosome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of HMGA2 that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that HMGA2 is involved in diet-induced obesity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized (provided by RefSeq, Jul 2008).				
Western Blotting	have been characterized (provided by RefSeq, Jul 2008). Mouse anti Human high mobility group protein HMGI-C antibody detects a band of 18 kDa. The antibody has been extensively validated for western blotting using whole cell				

Lysate Composition

400µg HepG2 lysate lyophilized in RIPA buffer

- $\textbf{Lysate Reconstitution} \ \ \textbf{-} \ \text{If using DDT reconstitute the lyophilized lysate with 190} \mu \ \text{DI H}_2\text{O}, \ \text{add 200} \mu \text{I } 2x \ \text{Laemmli}$ Sample Buffer and 10µl 2M DTT.
 - If using BME reconstitute the lyophilized lysate with 180 μ l DI H₂O, add 200 μ l 2x Laemmli Sample Buffer and 20µl BME.

Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

Storage

Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for

up to 3 months or at -80°C for longer term storage.

Guarantee	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 **Europe** Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351154:190315'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint