

Datasheet: VMA00470KT

Description:	PSMB9 ANTIBODY WITH CONTROL LYSATE
Specificity:	PSMB9
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Clone:	2B6
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 50% Glycerol
Immunogen	Recombinant protein fragment corresponding to amino acids 21-219 of human PSMB9 produced in <i>E. Coli</i>
External Database Links	UniProt: P28065 Related reagents Entrez Gene:

Synonyms	LMP2, PSMB6i, RING12
-----------------	----------------------

Specificity	<p>Mouse anti Human PSMB9 antibody recognizes the proteasome subunit beta type-9, also known as large multifunctional peptidase 2, low molecular mass protein 2, macropain chain 7, multicatalytic endopeptidase complex chain 7, proteasome chain 7, or proteasome subunit beta-1i.</p> <p>The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMB9 gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. PSMB9 is located in the class II region of the MHC (major histocompatibility complex). Expression of PSMB9 is induced by gamma interferon and PSMB9 product replaces catalytic subunit 1 (proteasome beta 6 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit (provided by RefSeq, Mar 2010).</p> <p>Mouse anti Human PSMB9 antibody detects a band of 23 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
--------------------	---

Western Blotting	Anti PSMB9 detects a band of approximately 23 kDa in HEK293 cell lysate
-------------------------	---

Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
-----------------------------	---

Lysate Composition	400µg HEK293 lysate lyophilized in RIPA buffer.
---------------------------	---

Lysate Reconstitution	<ul style="list-style-type: none">- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
------------------------------	--

Storage	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
----------------	--

Guarantee	As supplied, 12 months from date of despatch.
------------------	---

Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
-------------------------	--

Health And Safety Information	<p>Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</p>
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M338177:181217'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)