

Datasheet: VMA00488 BATCH NUMBER 160712

MOUSE ANTI PDIA1
PDIA1
Purified
PrecisionAb Monoclonal
lgG1
100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Recombinant protein of human PDIA1
External Database Links	UniProt: P07237 Related reagents

Entrez Gene:

5034 P4HB Related reagents

Synonyms	ERBA2L, PDI, PDIA1, PO4DB	
Specificity	Mouse anti Human PDIA1 antibody recognizes PDIA1, also kn hormone-binding protein, collagen prolyl 4-hydroxylase beta, glut transhydrogenase, p55, procollagen-proline, 2-oxoglutarate 4-dic polypeptide or thyroid hormone-binding protein p55.	tathione-insulin
	The P4HB gene encodes the beta subunit of prolyl 4-hydroxylase multifunctional enzyme that belongs to the protein disulfide isome present as a tetramer consisting of two alpha and two beta subunity involved in hydroxylation of prolyl residues in preprocollagen. The disulfide isomerase containing two thioredoxin domains that catabreakage and rearrangement of disulfide bonds. Other known furto act as a chaperone that inhibits aggregation of misfolded protein dependent manner, its ability to bind thyroid hormone, its role in of S-nitrosothiol-bound nitric oxide, and its function as a subunitariglyceride transfer protein complex (provided by RefSeq, Jul 20)	erase family. When nits, this enzyme is is enzyme is also a alyze the formation, nctions include its ability eins in a concentration-both the influx and efflux of the microsomal
	Mouse anti Human PDIA1 antibody detects a band of 57 kDa. The extensively validated for western blotting using whole cell lysates	•
Western Blotting	Anti PDIA1 detects a band of approximately 57 kDa in HeLa cell	lysate
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	
Guarantee	12 months from date of despatch	
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.	
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00488 Antibody (10040)	
Regulatory	For research purposes only	

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint