

Datasheet: VMA00490KT

Description:	HIF1AN ANTIBODY WITH CONTROL LYSATE
Specificity:	HIF1AN
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	lgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Full length human HIF1AN expressed in <i>E. Coli</i>
External Database Links	UniProt:

Q9NWT6 Related reagents

Entrez Gene:

55662 HIF1AN Related reagents

Synonyms	FIH1
Specificity	Mouse anti Human HIF1AN antibody recognizes HIF1AN also known as hypoxia-inducible factor 1-alpha inhibitor, factor inhibiting HIF-1, hypoxia-inducible factor asparagine hydroxylase or peptide-aspartate beta-dioxygenase.
	Mouse anti Human HIF1AN antibody detects a band of 42 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti HIF1AN detects a band of approximately 42 kDa in HEK293 cell lysate
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg HEK293 lysate lyophilized in RIPA buffer.
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT. If using BME reconstitute the lyophilized lysate with 180μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 20μl BME.
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50
 From

 Email: antibody_sales_us@bio-rad.com
 Email: antibody_sales_uk@bio-rad.com
 Email: antibody_sales_de@bio-rad.com
 Email: antibody_sales_de@bio-rad.com

15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351155:190315'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint