

Datasheet: VMA00498

Description:	MOUSE ANTI C-MYC
Specificity:	C-MYC
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG1
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	<i>E. coli</i> derived recombinant protein corresponding to amino acids 66-201 of human c-Myc
External Database Links	<p>UniProt: P01106 Related reagents</p> <p>Entrez Gene: 4609 MYC Related reagents</p>

Synonyms BHLHE39

Specificity **Mouse anti Human c-Myc antibody** recognizes the Myc proto-oncogene protein, also known as avian myelocytomatosis viral oncogene homolog, class E basic helix-loop-helix protein 39, myc-related translation/localization regulatory factor, transcription factor p64 or v-myc myelocytomatosis viral oncogene homolog.

The protein encoded by MYC gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of MYC have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma ([BL](#)). There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of MYC (provided by RefSeq, Jul 2008).

Mouse anti Human c-Myc antibody detects a band of 57 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Anti c-Myc detects a band of approximately 57 kDa in HeLa cell lysate

Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Guarantee As supplied, 12 months from date of despatch.

Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: Antibody (10040): <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M358540:190911'

Printed on 11 Oct 2019