

## Datasheet: VMA00498KT

<b>Description:</b>	C-MYC ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	C-MYC
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb Monoclonal
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	2 Westerns

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Mouse, Rat

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG - liquid

#### Preparation

20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant

#### Buffer Solution

Phosphate buffered saline

#### Preservative Stabilisers

0.09% Sodium Azide (NaN<sub>3</sub>)

**Immunogen** *E. coli* derived recombinant protein corresponding to amino acids 66-201 of human c-Myc

---

**External Database**

**Links**

**UniProt:**

[P01106](#) [Related reagents](#)

**Entrez Gene:**

[4609](#) MYC [Related reagents](#)

---

**Synonyms**

BHLHE39

---

**Specificity**

**Mouse anti Human c-Myc antibody** recognizes the Myc proto-oncogene protein, also known as avian myelocytomatosis viral oncogene homolog, class E basic helix-loop-helix protein 39, myc-related translation/localization regulatory factor, transcription factor p64 or v-myc myelocytomatosis viral oncogene homolog.

The protein encoded by MYC gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of MYC have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma ([BL](#)). There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of MYC (provided by RefSeq, Jul 2008).

Mouse anti Human c-Myc antibody detects a band of 57 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

---

**Western Blotting**

Anti c-Myc detects a band of approximately 57 kDa in HeLa cell lysate

---

**Instructions For Use**

Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

---

**Lysate Composition**

400µg HeLa lysate lyophilized in RIPA buffer.

---

**Lysate Reconstitution**

- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.

- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.

Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

---

**Storage**

Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.

---

**Guarantee** As supplied, 12 months from date of despatch.

---

**Acknowledgements** PrecisionAb™ is a trademark of Bio-Rad Laboratories.

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 #10561 available at:  
Antibody (10040): <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>  
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](http://bio-rad-antibodies.com)

'M350984:190315'

**Printed on 10 Feb 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)