

Datasheet: VMA00540KT

Description:	COMPLEMENT C3 ANTIBODY WITH CONTROL LYSATE
Specificity:	COMPLEMENT C3
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Clone:	AB03/3G11
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	<i>E. coli</i> -derived recombinant human complement C3 (amino acids 1200 - 1663)
External Database Links	<p>UniProt: P01024 Related reagents</p> <p>Entrez Gene: 718 C3 Related reagents</p>

Synonyms	CPAMD1
Specificity	<p>Mouse anti Human complement C3 antibody recognizes the complement C3, also known as C3 and PZP-like alpha-2-macroglobulin domain-containing protein 1, C3a anaphylatoxin, acylation-stimulating protein cleavage product, ASP or epididymis secretory sperm binding protein Li 62p.</p> <p>Complement component C3 plays a central role in the activation of complement system. Its activation is required for both classical and alternative complement activation pathways. People with C3 deficiency are susceptible to bacterial infection (provided by RefSeq, Feb 2009).</p> <p>Mouse anti Human complement C3 antibody detects bands of 115 and 68 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
Western Blotting	Anti complement C3 detects bands of approximately 68 kDa in C6 cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg C6 lysate lyophilized in RIPA buffer
Lysate Reconstitution	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
Storage	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	<p>Material Safety Datasheet documentation #10040 #10561 available at:</p> <p>Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</p> <p>Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</p>
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

'M345701:190125'

Printed on 15 Mar 2019