

## Datasheet: VMA00551KT

<b>Description:</b>	GSTM3 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	GSTM3
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb™ Monoclonal
<b>Clone:</b>	CPTC23
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	2 Westerns

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Immunogen</b>	Recombinant protein of human GSTM3
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P21266</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">2947</a> GSTM3   <a href="#">Related reagents</a></p>

<b>Synonyms</b>	GST5
<b>Specificity</b>	<p><b>Mouse anti Human GSTM3 antibody</b> recognizes the Glutathione S-transferase Mu 3, also known as GST class-mu 3, S-(hydroxyalkyl)glutathione lyase M3, brain GST, brain type mu-glutathione S-transferase or glutathione S-alkyltransferase M3.</p> <p>Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. GSTM3 gene encodes a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs. Mutations of this class mu gene have been linked with a slight increase in a number of cancers, likely due to exposure with environmental toxins. Alternative splicing results in multiple transcript variants (provided by RefSeq, Nov 2008).</p> <p>Mouse anti Human GSTM3 antibody detects a band of 26 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
<b>Western Blotting</b>	Anti GSTM3 detects a band of approximately 26 kDa in HeLa cell lysates
<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
<b>Lysate Composition</b>	400µg HeLa lysate lyophilized in RIPA buffer
<b>Lysate Reconstitution</b>	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
<b>Shelf Life</b>	As supplied, 12 months from date of despatch.
<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10040 #10561 available at:</p> <p>Antibody (10040): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a></p> <p>Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a></p>
<b>Regulatory</b>	For research purposes only

## Related Products

## Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

## Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M337202:181217'

**Printed on 02 Jan 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)