

Datasheet: VMA00564KT

Description:	CCAR2 ANTIBODY WITH CONTROL LYSATE
Specificity:	CCAR2
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Clone:	D02/1C5-4
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	<i>E. coli</i> -derived recombinant human CCAR2 (amino acids 5 - 366)
External Database Links	<p>UniProt: Q8N163 Related reagents</p> <p>Entrez Gene: 57805 KIAA1967 Related reagents</p>
Synonyms	DBC1

Specificity	<p>Mouse anti Human CCAR2 antibody recognizes the cell cycle and apoptosis regulator protein 2, also known as cell division cycle and apoptosis regulator protein 2, deleted in breast cancer 1 or p30 DBC protein.</p> <p>Mouse anti Human CCAR2 antibody detects a band of 130 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
Western Blotting	Anti CCAR2 detects a band of approximately 130 kDa in HEK293 cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg HEK293 lysate lyophilized in RIPA buffer
Lysate Reconstitution	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
Storage	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
Shelf Life	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	<p>Material Safety Datasheet documentation #10040 #10561 available at:</p> <p>Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</p> <p>Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</p>
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...)[HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M336482:181214'

Printed on 02 Jan 2019