Datasheet: VMA00584 BATCH NUMBER 161202

Description:	MOUSE ANTI PLC GAMMA-1
Specificity:	PLC GAMMA-1
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	1F1
lsotype:	lgG2a
Quantity:	100 µl

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Western Blotting	-			1/1000	
Target Species	The PrecisionAb label criteria within Bio-Rad ¹ how we validate our Pr use in a particular techni Further optimization may Human	s ongoing ecisionAl que this d	g antibod o range. \ oes not n	l y validation program Where this product ha ecessarily exclude its	nme. Click <u>here</u> to learn s not been tested for	
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity a reactivity is derived from personal communication further information.	and workir testing wi	thin our la	aboratories, peer-revie	wed publications or	
Product Form	Purified IgG - liquid					
Preparation	Mouse monoclonal antib	ody purifie	ed by affin	ity chromatography or	n Protein G from ascites	
Buffer Solution	HEPES buffered saline					
Preservative	0.09% Sodium Azide (Na	aN ₃)				

Approx. Protein ConcentrationsIgG concentration 0.5 mg/mlImmunogenE. coli-derived recombinant human PLC gamma-1External Database LinksUniProt: P19174 Related reagentsSynonymsPLC1SpecificityMouse anti Human PLC gamma-1 antibody recognizes the 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1, also known as PLC-148, phosphatidylinositol phosphodiesterase gama-1, also known as PLC-148, phosphatidylinositol also konverse second phosphodiesterase ga	Stabilisers	1% Bovine Serum Albumin <50% Glycerol		
External Databases UniProt: P19174 Related reagents Entrez Gone: 5335 Synonyms PLC1 Specificity Mouse anti Human PLC gamma-1 antibody recognizes the 1-phosphatidylinositol 4.5-bisphosphate phosphodiesterase gamma-1, also known as PLC-148, phosphatidylinositol phospholipase C or phosphotiositidase C. The protein encoded by PLCG1 gene catalyzes the formation of inositol 1,4,5- trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein hasebeen shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)- activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008). Western Blotting Anti PLC gamma-1 detects a band of approximately 150 kDa in Jurkat cell lysates Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)		IgG concentration 0.5 mg/ml		
Links UniProt: P19174 Related reagents Entrez Gone: 5335 5335 PLCG1 Related reagents Synonyms PLC1 Specificity Mouse anti Human PLC gamma-1 antibody recognizes the 1-phosphatidylinositol 4,5-bisphosphate phospholipase C or phosphoinositidase C. The protein encoded by PLCG1 gene catalyzes the formation of inositol 1,4,5- trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)- activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008). Mouse anti Human PLC gamma-1 antibody detects a band of 150 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #20359 available at: https://www bio-rad-antibodies.com/SDS/WA00584 Antibody	Immunogen	<i>E. coli</i> -derived recombinant human PLC gamma-1		
Specificity Mouse anti Human PLC gamma-1 antibody recognizes the 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1, also known as PLC-148, phosphatidylinositol phosphodipase C or phosphonositidase C. The protein encoded by PLCG1 gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008). Western Blotting Anti PLC gamma-1 detects a band of approximately 150 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)		P19174 Related reagents Entrez Gene:		
4.5-bisphosphate phospholiesterase gamma-1, also known as PLC-148, phosphatidylinositol phospholipase C or phosphoinositidase C. The protein encoded by PLCG1 gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008). Mouse anti Human PLC gamma-1 antibody detects a band of 150 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584_Antibody (20359)	Synonyms	PLC1		
trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)- activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008).Western BlottingAnti PLC gamma-1 detects a band of approximately 150 kDa in Jurkat cell lysatesStorageStore undiluted at -20°C, avoiding repeated freeze thaw cycles.Guarantee12 months from date of despatchAcknowledgementsPrecisionAb is a trademark of Bio-Rad Laboratories.Health And Safety InformationMaterial Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)	Specificity	4,5-bisphosphate phosphodiesterase gamma-1, also known as phosphatidylinositol phospholipase C or phosphoinositidase C.	PLC-148,	
been extensively validated for western blotting using whole cell lysates.Western BlottingAnti PLC gamma-1 detects a band of approximately 150 kDa in Jurkat cell lysatesStorageStore undiluted at -20°C, avoiding repeated freeze thaw cycles.Guarantee12 months from date of despatchAcknowledgementsPrecisionAb is a trademark of Bio-Rad Laboratories.Health And Safety InformationMaterial Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)		trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)- activated tyrosine kinase. Two transcript variants encoding different isoforms have been		
StorageStore undiluted at -20°C, avoiding repeated freeze thaw cycles.Guarantee12 months from date of despatchAcknowledgementsPrecisionAb is a trademark of Bio-Rad Laboratories.Health And Safety InformationMaterial Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)		č		
Guarantee12 months from date of despatchAcknowledgementsPrecisionAb is a trademark of Bio-Rad Laboratories.Health And Safety InformationMaterial Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)	Western Blotting	Anti PLC gamma-1 detects a band of approximately 150 kDa in	n Jurkat cell lysates	
Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #20359 available at: Information https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)	Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles		
Health And Safety Material Safety Datasheet documentation #20359 available at: Information https://www.bio-rad-antibodies.com/SDS/VMA00584 Antibody (20359)	Guarantee	12 months from date of despatch		
Information <u>https://www.bio-rad-antibodies.com/SDS/VMA00584</u> Antibody (20359)	Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.		
Regulatory For research purposes only	-	https://www.bio-rad-antibodies.com/SDS/VMA00584		
	Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bi	o-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M370298:200529'

Printed on 01 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint