

Datasheet: VMA00584

Description:	MOUSE ANTI PLC GAMMA-1	
Specificity:	PLC GAMMA-1	
Format:	Purified	
Product Type:	PrecisionAb Monoclonal	
Clone:	1F1	
Isotype:	IgG2a	
Quantity:	100 μΙ	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Learn about how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody purified by affinity chromatography on Protein G from ascites.
Buffer Solution	HEPES buffered saline.
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin <50% Glycerol

Approx. Protein Concentrations	IgG concentration 0.5 mg/ml.
Immunogen	E. coli-derived recombinant human PLC gamma-1
External Database Links	UniProt: P19174 Related reagents Entrez Gene: 5335 PLCG1 Related reagents
Synonyms	PLC1
Specificity	Mouse anti Human PLC gamma-1 antibody recognizes the 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1, also known as PLC-148, phosphatidylinositol phospholipase C or phosphoinositidase C. The protein encoded by PLCG1 gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008). Mouse anti Human PLC gamma-1 antibody detects a band of 150 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti PLC gamma-1 detects a band of approximately 150 kDa in Jurkat cell lysates.
Storage	This product is shipped at ambient temperature. Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Guarantee	12 months from date of despatch.
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #20359 available at: https://www.bio-rad-antibodies.com/SDS/VMA00584
Regulatory	For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M443639:250709'

Printed on 09 Jul 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint