

Datasheet: VMA00584KT

Description:	PLC GAMMA-1 ANTIBODY WITH CONTROL LYSATE
Specificity:	PLC GAMMA-1
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Isotype:	lgG2a
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human	
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.	
Product Form	Purified IgG - liquid	
Preparation	20μl Mouse monoclonal antibody purified by affinity chromatography Protein G from ascites	
Buffer Solution	HEPES buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin <50% Glycerol	
Immunogen	E. coli-derived recombinant human PLC gamma-1	
External Database Links	UniProt: P19174 Related reagents Entrez Gene: 5335 PLCG1 Related reagents	
Synonyms	PLC1	

Specificity	Mouse anti Human PLC gamma-1 antibody recognizes the 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1, also known as PLC-148, phosphatidylinositol phospholipase C or phosphoinositidase C.
	The protein encoded by PLCG1 gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for PLCG1 (provided by RefSeq, Jul 2008).
	Mouse anti Human PLC gamma-1 antibody detects a band of 150 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti PLC gamma-1 detects a band of approximately 150 kDa in Jurkat cell lysates
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg Jurkat lysate lyophilized in RIPA buffer
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT. If using BME reconstitute the lyophilized lysate with 180μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 20μl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #20359 #10561 available at: Antibody (20359): https://www.bio-rad-antibodies.com/uploads/MSDS/20359.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com 'M338031:181217'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint