

## Datasheet: VMA00599

<b>Description:</b>	HUMAN ANTI UBE2D2
<b>Specificity:</b>	UBE2D2
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb™ Monoclonal
<b>Isotype:</b>	HuCAL Fab bivalent
<b>Quantity:</b>	100 µl

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Human monoclonal antibody purified by metal chelate affinity chromatography from <i>E. coli</i> extract
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Immunogen</b>	<i>E. coli</i> -derived full length recombinant human UBE2D2
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P62837</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">7322</a>   UBE2D2   <a href="#">Related reagents</a></p>
<b>Synonyms</b>	UBC4, UBC5B, UBCH4, UBCH5B

**Specificity** **Human anti Human UBE2D2 antibody** recognizes the ubiquitin-conjugating enzyme E2 D2, also known as p53-regulated ubiquitin-conjugating enzyme 1, ubiquitin carrier protein D2, ubiquitin-conjugating enzyme E2-17 kDa 2 or Ubiquitin-protein ligase D2.

Regulated degradation of misfolded, damaged or short-lived proteins in eukaryotes occurs via the ubiquitin (Ub)-proteasome system (UPS). An integral part of the UPS system is the ubiquitination of target proteins and covalent linkage of Ub-containing proteins to form polymeric chains, marking them as targets for 26S proteasome-mediated degradation. Ubiquitination of proteins is mediated by a cascade of enzymes which includes E1 (ubiquitin activating), E2 (ubiquitin conjugating), and E3 (ubiquitin ligases) enzymes. UBE2D2 encodes a member of the E2 enzyme family. Substrates of this enzyme include the tumor suppressor protein p53 and peroxisomal biogenesis factor 5 (PEX5). Alternative splicing results in multiple transcript variants of UBE2D2 (provided by RefSeq, May 2013).

Human anti Human UBE2D2 antibody detects a band of 18 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

---

**Western Blotting** Human anti UBE2D2 antibody detects a band of approximately 18 kDa in NIH/3T3 cell lysates

---

**Instructions For Use** Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

---

**Storage** Store undiluted at -20°C, avoiding repeated freeze thaw cycles

---

**Guarantee** As supplied, 12 months from date of despatch

---

**Acknowledgements** PrecisionAb™ is a trademark of Bio-Rad Laboratories

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: Antibody (10040): <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Human IgG F(ab')<sub>2</sub> (STAR126...) [HRP](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M338897:181217'

Printed on 20 May 2019