

Datasheet: VMA00604KT

Description:	APC2 ANTIBODY WITH CONTROL LYSATE
Specificity:	APC2
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Clone:	OTI1B11
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Human			
Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species.			
Purified IgG - liquid			
20μl Mouse monoclonal antibody purified by affinity chromatograpy on Protein G from ascites			
Phosphate buffered saline			
0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin <50% Glycerol			
HEK293-derived full length recombinant human APC2			
UniProt: Q9UJX6 Related reagents Entrez Gene:			
	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. Purified IgG - liquid 20µl Mouse monoclonal antibody purified by affinity chromatograpy on Protein G from a Phosphate buffered saline 0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin <50% Glycerol HEK293-derived full length recombinant human APC2 UniProt: Q9UJX6 Related reagents		

Specificity Mouse anti Human APC2 antibody recognizes the anaphase promoting complex subunit 2 also known as cyclosome subunit 2. A large protein complex, termed the anaphase-promoting complex (APC) or the cyclosome, promotes metaphase-anaphase transition by ubiquitinating its specific substrates such as mitotic cyclins and anaphase inhibitor, which are subsequently degraded by the 26S proteasome. Biochemical studies have shown that the vertebrate APC contains eight subunits. The composition of the APC is highly conserved in organisms from yeast to humans. The product of this gene is a component of the complex and shares sequence similarity with a recently identified family of proteins called cullins, which may also be involved in ubiquitin-mediated degradation (provided by RefSeq. Jul 2008). Mouse anti Human APC2 antibody detects a band of 100kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting	Synonyms	APC2, KIAA1406
promotes metaphase-anaphase transition by ubiquitinating its specific substrates such as mitotic cyclins and anaphase inhibitor, which are subsequently degraded by the 26S proteasome. Biochemical studies have shown that the verterbate APC contains eight submits. The composition of the APC is highly conserved in organisms from yeast to humans. The product of this gene is a component of the complex and shares sequence similarity with a recently identified family of proteins called cullins, which may also be involved in ubiquitin-mediated degradation (provided by RefSeq, Jul 2008). Mouse anti Human APC2 antibody detects a band of 100kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Mouse anti APC2 antibody detects a band of approximately 100 kDa in HEK293 cell lysates Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch Acknowledgements Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Specificity	
walidated for western blotting using whole cell lysates. Western Blotting Mouse anti APC2 antibody detects a band of approximately 100 kDa in HEK293 cell lysates Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at:		promotes metaphase-anaphase transition by ubiquitinating its specific substrates such as mitotic cyclins and anaphase inhibitor, which are subsequently degraded by the 26S proteasome. Biochemical studies have shown that the vertebrate APC contains eight subunits. The composition of the APC is highly conserved in organisms from yeast to humans. The product of this gene is a component of the complex and shares sequence similarity with a recently identified family of proteins called cullins, which may also be involved in ubiquitin-mediated degradation (provided by
Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition 400μg HEK293 lysate lyophilized in RIPA buffer Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 20μl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch PrecisionAb™ is a trademark of Bio-Rad Laboratories Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf		
Lysate Composition Lysate Composition Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10581.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10581.pdf	Western Blotting	Mouse anti APC2 antibody detects a band of approximately 100 kDa in HEK293 cell lysates
Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Instructions For Use	•
Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Lysate Composition	400μg HEK293 lysate lyophilized in RIPA buffer
refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Lysate Reconstitution	Buffer and 10 μ l 2M DTT. - If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME.
Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf		
months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf		
Health And Safety Information Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Guarantee	As supplied, 12 months from date of despatch
Information Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Regulatory For research purposes only	-	Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf
	Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M347894:190214'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint