

Datasheet: VMA00627

Description:	MOUSE ANTI UPF1
Specificity:	UPF1
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	AB03/4D5
lsotype:	lgG2a
Quantity:	100 µl

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
	rad-antibodies.com/prote	Yes	No	Not Determined	Suggested Dilution	
	Immunoprecipitation	163	NO	Not Determined	Suggested Dilution	
	Western Blotting	•			1/1000	
	The PrecisionAb label	is reserve	d for an	tibodies that meet the		
	criteria within Bio-Rad				•	
	we validate our Precisi	-	-			
	particular technique this			•		
	optimization may be req			•	ch procedures. Further	
Target Species	Human					
Species Cross	Reacts with: Mouse, Ra	t				
Reactivity	N.B. Antibody reactivity	and worki	ng conditi	ions may vary between	species. Cross	
	reactivity is derived from personal communicatior further information.	-		•	•	
Product Form	Purified IgG - liquid					
Preparation	Mouse monoclonal antib culture supernatant.	ody purifi	ed by affir	nity chromatograpy on	Protein G from tissue	
Buffer Solution	Phosphate buffered sali	ne.				
Preservative	0.09% Sodium Azide (N	aN ₃).				

Stabilisers

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml.
Immunogen	<i>E. coli</i> -derived recombinant human UPF1 (aa 630-1129)
External Database Links	UniProt: <u>Q92900</u> <u>Related reagents</u> Entrez Gene:
	5976 UPF1 Related reagents
Synonyms	KIAA0221, RENT1
Specificity	Mouse anti Human UPF1 antibody recognizes UPF1, also known as RENT1, NORF1, regulator of nonsense transcripts 1 or delta helicase.
	This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located only in the cytoplasm. When translation ends, it interacts with the protein that is a functional homolog of yeast Upf2p to trigger mRNA decapping. Use of multiple polyadenylation sites has been noted for this gene. Alternative splicing results in multiple transcript variants (provided by RefSeq, Jul 2014). Mouse anti Human UPF1 antibody detects a band of 143 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Mouse anti UPF1 antibody detects a band of approximately 143 kDa in HEK293 cell lysates.
Storage	This product is shipped at ambient temperature. Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Guarantee	12 months from date of despatch.
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00627
Regulatory	For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...)<u>HRP</u> Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M443670:250709'

Printed on 09 Jul 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint