

## Datasheet: VMA00627KT

| Description:         | UPF1 ANTIBODY WITH CONTROL LYSATE |
|----------------------|-----------------------------------|
| Specificity:         | UPF1                              |
| Format:              | Purified                          |
| <b>Product Type:</b> | PrecisionAb™ Monoclonal           |
| Clone:               | AB03/4D5                          |
| Isotype:             | lgG2a                             |
| Quantity:            | 2 Westerns                        |

### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                  | Yes | No | Not Determined | Suggested Dilution |
|------------------|-----|----|----------------|--------------------|
| Western Blotting | •   |    |                | 1/1000             |

**PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.** 

| Target Species              | Human  |  |  |
|-----------------------------|--|--|--|
| Species Cross<br>Reactivity | Reacts with: Mouse, Rat  N.B. Antibody reactivity and working conditions may vary between species.             |  |  |
| Product Form                | Purified IgG - liquid  |  |  |
| Preparation                 | 20µl Mouse monoclonal antibody purified by affinity chromatograpy on Protein G from tissue culture supernatant |  |  |
| Buffer Solution             | Phosphate buffered saline  |  |  |
| Preservative<br>Stabilisers | 0.09% Sodium Azide (NaN <sub>3</sub> )   |  |  |
| Immunogen                   | E. coli-derived recombinant human UPF1 (aa 630-1129)   |  |  |
| External Database<br>Links  | UniProt:  Q92900 Related reagents  |  |  |
|                             | Entrez Gene: <u>5976</u> UPF1 Related reagents   |  |  |

| Synonyms                         | KIAA0221, RENT1  |
|----------------------------------|--|
| Specificity                      | Mouse anti Human UPF1 antibody recognizes UPF1, also known as RENT1, NORF1, regulator of nonsense transcripts 1 or delta helicase.   |
|                                  | This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located only in the cytoplasm. When translation ends, it interacts with the protein that is a functional homolog of yeast Upf2p to trigger mRNA decapping. Use of multiple polyadenylation sites has been noted for this gene. Alternative splicing results in multiple transcript variants (provided by RefSeq, Jul 2014). |
|                                  | Mouse anti Human UPF1 antibody detects a band of 143 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.   |
| Western Blotting                 | Mouse anti UPF1 antibody detects a band of approximately 143 kDa in HEK293 cell lysates  |
| Instructions For Use             | Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.  |
| Lysate Composition               | 400μg HEK293 lysate lyophilized in RIPA buffer   |
| Lysate Reconstitution            | <ul> <li>If using DDT reconstitute the lyophilized lysate with 190μl DI H<sub>2</sub>O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT.</li> <li>If using BME reconstitute the lyophilized lysate with 180μl DI H<sub>2</sub>O, add 200μl 2x Laemmli Sample Buffer and 20μl BME.</li> <li>Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please</li> </ul>  |
|                                  | refer to the PrecisionAb western blotting protocol.  |
| Storage                          | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.  |
|                                  | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.   |
| Guarantee                        | As supplied, 12 months from date of despatch   |
| Acknowledgements                 | PrecisionAb™ is a trademark of Bio-Rad Laboratories  |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a> Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a>   |
| Regulatory                       | For research purposes only   |

## **Related Products**

# **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

**Recommended Negative Controls** 

### MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

'M338924:181217'

#### Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint