

## Datasheet: VMA00657KT

<b>Description:</b>	p27/Kip1 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	p27/Kip1
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb™ Monoclonal
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	2 Westerns

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Immunogen</b>	Full length recombinant human p27/Kip1
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P46527</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">1027</a>   CDKN1B   <a href="#">Related reagents</a></p>
<b>Synonyms</b>	KIP1

<b>Specificity</b>	<p><b>Mouse anti Human p27/Kip1 antibody</b> recognizes cyclin-dependent kinase inhibitor 1, also known as CDK-interacting protein 1, KIP1, BWS, P27KIP1 or wild-type p53-activated fragment 1.</p> <p>CDKN1A encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of CDKN1A is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen (PCNA), a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation. Multiple alternatively spliced variants have been found for CDKN1A (provided by RefSeq, Nov 2010).</p> <p>Mouse anti Human p27/Kip1 antibody detects a band of 27 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
<b>Western Blotting</b>	Mouse anti p27/Kip1 antibody detects a band of approximately 27 kDa in C6 cell lysates
<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
<b>Lysate Composition</b>	400µg C6 lysate lyophilized in RIPA buffer
<b>Lysate Reconstitution</b>	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
<b>Guarantee</b>	As supplied, 12 months from date of despatch
<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories
<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10040 #10561 available at:</p> <p>Antibody (10040): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a></p> <p>Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a></p>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

### Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M337860:181217'

**Printed on 16 Mar 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)