

Datasheet: VMA00820 BATCH NUMBER 64619670

Specificity: hnRNP C Format: Purified Product Type: PrecisionAb Monoclo	C
Product Type: PrecisionAb Monoclo	
	nal
Clone: AB05/4F12	
lsotype: IgG1	
Quantity : 100 μl	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunoprecipitation	•			
Western Blotting	•			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - Liquid
Preparation	Mouse monoclonal antibody affinity purified on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	E. coli-derived recombinant protein of amino acids 1-306 of hu	man hnRNP C
External Database Links	UniProt: P07910 Related reagents Entrez Gene: 3183 HNRNPC Related reagents	
Synonyms	HNRPC	
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cel myeloma cell line	lls of the mouse SP2/0
Specificity	Mouse anti hnRNP C antibody recognizes heterogeneous nu C1/C2, also known as HNRPC.	ıclear ribonucleoproteins
	hnRNP C is a ubiquitously expressed nuclear RNA-binding pro- all nascent mRNA transcripts (<u>Zarnack et al. 2013</u>). This protein RNA maturation including pre-mRNA splicing, mRNA stability, modulation (<u>Nasrin et al. 2014</u>), and one of its main functions at the human transcriptome from aberrant Alu exonization (<u>Zarnach</u> hnRNP C plays roles in cell proliferation and differentiation. Du cell cycle, hnRNP C translocates from the nucleus to the cytop affect translation and regulate mitosis (<u>Yang et al. 2013</u>).	in is involved with stages of and translational appears to be protection of ack et al. 2013). In addition, uring the G2/M phase of the
Western Blotting	all nascent mRNA transcripts (<u>Zarnack et al. 2013</u>). This protein RNA maturation including pre-mRNA splicing, mRNA stability, modulation (<u>Nasrin et al. 2014</u>), and one of its main functions at the human transcriptome from aberrant Alu exonization (<u>Zarnatharnach in RNP C plays roles in cell proliferation and differentiation. Du cell cycle, hnRNP C translocates from the nucleus to the cytop</u>	in is involved with stages of and translational appears to be protection of tok et al. 2013). In addition, uring the G2/M phase of the plasm where it is able to
Western Blotting Storage	all nascent mRNA transcripts (Zarnack et al. 2013). This protein RNA maturation including pre-mRNA splicing, mRNA stability, modulation (Nasrin et al. 2014), and one of its main functions at the human transcriptome from aberrant Alu exonization (ZarnathnRNP C plays roles in cell proliferation and differentiation. Ducell cycle, hnRNP C translocates from the nucleus to the cytopaffect translation and regulate mitosis (Yang et al. 2013). Mouse anti hnRNP C antibody detects a band of approximately	in is involved with stages of and translational appears to be protection of tok et al. 2013). In addition, uring the G2/M phase of the blasm where it is able to
	all nascent mRNA transcripts (<u>Zarnack et al. 2013</u>). This protein RNA maturation including pre-mRNA splicing, mRNA stability, modulation (<u>Nasrin et al. 2014</u>), and one of its main functions at the human transcriptome from aberrant Alu exonization (<u>Zarnatharnach nastation</u>). Ducell cycle, hnRNP C translocates from the nucleus to the cytopaffect translation and regulate mitosis (<u>Yang et al. 2013</u>). Mouse anti hnRNP C antibody detects a band of approximately lysates	in is involved with stages of and translational appears to be protection of tok et al. 2013). In addition, uring the G2/M phase of the blasm where it is able to
Storage	all nascent mRNA transcripts (Zarnack et al. 2013). This protein RNA maturation including pre-mRNA splicing, mRNA stability, modulation (Nasrin et al. 2014), and one of its main functions at the human transcriptome from aberrant Alu exonization (Zarnath hnRNP C plays roles in cell proliferation and differentiation. Ducell cycle, hnRNP C translocates from the nucleus to the cytopaffect translation and regulate mitosis (Yang et al. 2013). Mouse anti hnRNP C antibody detects a band of approximately lysates Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	in is involved with stages of and translational appears to be protection of tok et al. 2013). In addition, uring the G2/M phase of the blasm where it is able to
Storage Guarantee	all nascent mRNA transcripts (Zarnack et al. 2013). This protein RNA maturation including pre-mRNA splicing, mRNA stability, modulation (Nasrin et al. 2014), and one of its main functions at the human transcriptome from aberrant Alu exonization (Zarnath hnRNP C plays roles in cell proliferation and differentiation. Ducell cycle, hnRNP C translocates from the nucleus to the cytopaffect translation and regulate mitosis (Yang et al. 2013). Mouse anti hnRNP C antibody detects a band of approximately lysates Store undiluted at -20°C, avoiding repeated freeze thaw cycles 12 months from date of despatch	in is involved with stages of and translational appears to be protection of tok et al. 2013). In addition, uring the G2/M phase of the plasm where it is able to

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M429607:240410'

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint