

Datasheet: VMA00870

Description:	MOUSE ANTI COFILIN 1
Specificity:	COFILIN 1
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	25C1-E3-D3
lsotype:	lgG2a
Quantity:	100 µl

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .								
		Yes	No	Not Determined	Suggested Dilution				
	Western Blotting	-			1/1000				
	criteria within Bio-Rad how we validate our Pr	s ongoing ecisionAl que this d	g antibod b range. \ oes not n	l y validation progra n Where this product ha ecessarily exclude its	e defined performance nme. Click <u>here</u> to learn as not been tested for use in such procedures.				
Target Species	Human								
Product Form	Purified IgG - liquid								
Preparation	Mouse monoclonal antibody purified by affinity chromatography on Protein A from tissue culture supernatant								
Buffer Solution	Phosphate buffered salir	ie							
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)								
Approx. Protein Concentrations	IgG concentration 0.5 m	g/ml							
Immunogen	E. coli-derived recombin	ant humar	n cofilin 1	(aa 1-166)					

External Database Links	UniProt: <u>P23528</u> <u>Related reagents</u> Entrez Gene: <u>1072</u> CFL1 <u>Related reagents</u>					
Synonyms	CFL					
Specificity	Mouse anti Human cofilin 1 antibody recognizes CFL1, also known as HEL-S-15.					
	 The protein encoded by this gene can polymerize and depolymerize F-actin and G-actin in a pH-dependent manner. Increased phosphorylation of this protein by LIM kinase aids in Rho-induced reorganization of the actin cytoskeleton. Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner. It is involved in the translocation of actin-cofilin complex from cytoplasm to nucleus (supplied by OMIM, Apr 2004). Mouse anti Human cofilin 1 antibody detects a band of 18 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. 					
Western Blotting	Mouse anti cofilin 1 antibody detects a band of approximately 18 kDa in HEK293 cells					
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles					
Guarantee	12 months from date of despatch					
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories					
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00870 10040					
Regulatory	For research purposes only					

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M394561:220215'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint