

Datasheet: VMA00922

Description:	MOUSE ANTI SRC
Specificity:	SRC
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	ST03/3E9
lsotype:	lgG2a
Quantity:	100 µl

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal							
	communications from the originators. Please refer to references indicated for fur							
	information. For general	information. For general protocol recommendations, please visit www.bio-						
	rad-antibodies.com/protocols.							
		Yes	No	Not Determined	Suggested Dilution			
	Western Blotting	-			1/1000			
	Immunofluorescence	-						
Target Species	how we validate our P	recisionA	b range. does not r	Where this product ha necessarily exclude its	nme. Click <u>here</u> to learn s not been tested for use in such procedures.			
Species Cross Reactivity	Reacts with: Rat, Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.							
Product Form	Purified IgG - Liquid							
Preparation	Mouse monoclonal antibody affinity purified on Protein G from tissue culture supernatant							
Buffer Solution	Phosphate buffered sali	ne						
Preservative	0.09% Sodium Azide							

Stabilisers				
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	E. coli-derived recombinant protein of amino acids 351-543 of human SRC			
External Database Links	UniProt: P12931 Related reagents Entrez Gene: <u>6714</u> SRC <u>Related reagents</u>			
Synonyms	SRC1			
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line			
Specificity	Mouse anti SRC antibody recognizes proto-oncogene tyrosine-protein kinase Src, also known as Src1, p60-Src and c-Src. c-Src is a ubiquitously-expressed cytoplasmic tyrosine kinase, important for regulating diverse cellular functions (Honda et al. 2018). c-Src was the first oncogene to be identified, overexpressed and abnormally activated in cancer cells, and is the cellular counterpart of v-Src, a cancer-causing gene found in the Rous sarcoma virus (Hunter and Sefton 1980). Src functions as a complex molecular switch, which can self-modulate its activity in response to specific signals. This is achieved by a reciprocally regulatory mechanism involving Tyr416 and Tyr527, changing the tertiary structure of Src and controlling access to its active site (Knighton et al. 1991).			
Western Blotting	Mouse anti SRC detects a band of approximately 58 kDa in A431 cell lysates			
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles			
Guarantee	12 months from date of despatch			
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00922			
	Antibody (10040)			

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-r	ad.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419837:230622'

Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint