

Datasheet: VMA00928

Description:	MOUSE ANTI BAG2	
Specificity:	BAG2	
Format:	Purified	
<b>Product Type:</b>	PrecisionAb Monoclonal	
Clone:	AB01/3G9	
Isotype:	lgG1	
Quantity:	100 μΙ	

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click <a href="here">here</a> to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - Liquid
Preparation	Mouse monoclonal antibody affinity purified on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	E. coli-derived recombinant protein of amino acids 1-211 of human BAG2

### **External Database** Links

**UniProt:** 

O95816 Related reagents

**Entrez Gene:** 

9532 BAG2 Related reagents

**Fusion Partners** 

Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line

**Specificity** 

Mouse anti BAG2 antibody recognizes BAG family molecular chaperone regulator 2, also known as Bcl-2-associated athanogene 2.

BAG2 is a member of the BAG family. These proteins interact with the molecular chapherone HSP70, dictating HSP70 interactions with client proteins and apoptosis. BAG2 also interacts with a range of other molecules, regulating various cellular functions (Qin et al. 2016). BAG2 phosphorylation by MAPKAP kinase 2 is part of the p38 signaling pathway in response to extracellular stress (Ueda et al. 2004). BAG2 is induced by proteasome inhibitors and appears to promote apoptosis in cancer cells (Wang et al. 2008). BAG2 has also been implicated in Parkinson's disease (Fukuzono et al. 2016) and Alzheimer's disease (de Paula et al. 2016). BAG2 may be involved in simian virus 40 infection (Dupzyk and Tsai, 2018).

Western Blotting	Mouse anti BAG2 detects a band of approximately 24 kDa in HeLa cell lysates
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles
Guarantee	12 months from date of despatch
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/VMA00928">https://www.bio-rad-antibodies.com/SDS/VMA00928</a> Antibody (10040)
Regulatory	For research purposes only

# Related Products

### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

# Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint