

# Datasheet: VMA00936 BATCH NUMBER 100005658

Description:	MOUSE ANTI NCF2	
Specificity:	NCF2	
Format:	Purified	
<b>Product Type:</b>	PrecisionAb Monoclonal	
Clone:	H05/3G6	
Isotype:	lgG1	
Quantity:	100 μΙ	

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Immunoprecipitation	•			
Western Blotting	-			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click <a href="here">here</a> to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - Liquid
Preparation	Mouse monoclonal antibody affinity purified on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	E. coli-derived recombinant protein of amino acids 1-526 of human NCF2
External Database Links	UniProt: P19878 Related reagents  Entrez Gene: 4688 NCF2 Related reagents
Synonyms	NOXA2, P67PHOX
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line
Specificity	Mouse anti NCF2 antibody recognizes neutrophil cytosol factor 2, also known as NOXA2 and P67PHOX.  NCF2 is a cytosolic component of NADPH oxidase, necessary for phagocyte reactive oxygen species (ROS) production, phagocytic microbicidal activity and innate immunity (Zhang et al. 2018). NCF2 is upregulated in response to p53 induction, and NCF2
	knockdown results in reduced production of ROS and stimulates cell death ( <a href="Italiano et al.2012">Italiano et al.2012</a> ). A variant of the gene encoding NCF2 causes altered protein binding and is associated with very early inflammatory bowel disease ( <a href="Muise et al.2012">Muise et al.2012</a> ). Mutations in NCF2 can result in granulomatous disease, an immunodeficiency characterized by recurrent infections ( <a href="Tarazona-Santos et al.2013">Tarazona-Santos et al.2013</a> ). Pattern of NCF2 expression has been suggested as a potential diagnostic tool in cervical cancer ( <a href="Lomnytska et al.2011">Lomnytska et al.2011</a> ).
Western Blotting	knockdown results in reduced production of ROS and stimulates cell death ( <u>Italiano et al. 2012</u> ). A variant of the gene encoding NCF2 causes altered protein binding and is associated with very early inflammatory bowel disease ( <u>Muise et al. 2012</u> ). Mutations in NCF2 can result in granulomatous disease, an immunodeficiency characterized by recurrent infections ( <u>Tarazona-Santos et al. 2013</u> ). Pattern of NCF2 expression has been
Western Blotting Storage	knockdown results in reduced production of ROS and stimulates cell death ( <u>Italiano et al. 2012</u> ). A variant of the gene encoding NCF2 causes altered protein binding and is associated with very early inflammatory bowel disease ( <u>Muise et al. 2012</u> ). Mutations in NCF2 can result in granulomatous disease, an immunodeficiency characterized by recurrent infections ( <u>Tarazona-Santos et al. 2013</u> ). Pattern of NCF2 expression has been suggested as a potential diagnostic tool in cervical cancer ( <u>Lomnytska et al. 2011</u> ).
	knockdown results in reduced production of ROS and stimulates cell death ( <u>Italiano et al. 2012</u> ). A variant of the gene encoding NCF2 causes altered protein binding and is associated with very early inflammatory bowel disease ( <u>Muise et al. 2012</u> ). Mutations in NCF2 can result in granulomatous disease, an immunodeficiency characterized by recurrent infections ( <u>Tarazona-Santos et al. 2013</u> ). Pattern of NCF2 expression has been suggested as a potential diagnostic tool in cervical cancer ( <u>Lomnytska et al. 2011</u> ).  Mouse anti NCF2 detects a band of approximately 67 kDa in RAW264.7 cell lysates
Storage	knockdown results in reduced production of ROS and stimulates cell death ( <u>Italiano et al. 2012</u> ). A variant of the gene encoding NCF2 causes altered protein binding and is associated with very early inflammatory bowel disease ( <u>Muise et al. 2012</u> ). Mutations in NCF2 can result in granulomatous disease, an immunodeficiency characterized by recurrent infections ( <u>Tarazona-Santos et al. 2013</u> ). Pattern of NCF2 expression has been suggested as a potential diagnostic tool in cervical cancer ( <u>Lomnytska et al. 2011</u> ).  Mouse anti NCF2 detects a band of approximately 67 kDa in RAW264.7 cell lysates  Store undiluted at -20°C, avoiding repeated freeze thaw cycles
Storage Guarantee	knockdown results in reduced production of ROS and stimulates cell death (

# **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M387461:210629'

### Printed on 10 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint