

Datasheet: VPA00114

Description:	GOAT ANTI BAG3
Specificity:	BAG3
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Goat polyclonal antibody purified by affinity chromatography.
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.02% Sodium Azide (NaN ₃) 0.5 % BSA
Immunogen	Peptide with the sequence C-SSMTDTPGNPAAP, from the C terminus of the protein sequence.
External Database Links	<p>UniProt: O95817 Related reagents</p> <p>Entrez Gene: 9531 BAG3 Related reagents</p>
Synonyms	BIS

Specificity **Goat anti Human BAG3 antibody** recognizes the BAG family molecular chaperone regulator 3 (BAG3), also known as Bcl-2-associated athanogene 3, Bcl-2-binding protein Bis and docking

protein CAIR-1.

BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. BAG3 contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner (provided by RefSeq, Jul 2008).

Goat anti Human BAG3 antibody detects a band of 85 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti BAG3 detects a band of approximately 85 kDa in MCF7 cell lysates.
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Guarantee	12 months from date of despatch
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10058 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M370506:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)