

Datasheet: VPA00114KT

Description:	BAG3 ANTIBODY WITH CONTROL LYSATE
Specificity:	BAG3
Format:	Purified
<b>Product Type:</b>	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

#### **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

**PrecisionAb antibodies have been extensively** <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human		
Product Form	Purified IgG - liquid		
Preparation	20μl Goat polyclonal antibody purified by affinity chromatography		
Buffer Solution	TRIS buffered saline		
Preservative Stabilisers	0.02% Sodium Azide (NaN <sub>3</sub> ) 0.5 % BSA		
Immunogen	Peptide with the sequence C-SSMTDTPGNPAAP, from the C terminus of the protein sequence.		
External Database Links	UniProt: O95817 Related reagents		

## **Entrez Gene:**

9531 BAG3 Related reagents

Synonyms	BIS
Specificity	<b>Goat anti Human BAG3 antibody</b> recognizes the BAG family molecular chaperone regulator 3 (BAG3), also known as Bcl-2-associated athanogene 3, Bcl-2-binding protein Bis and docking protein CAIR-1.
	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. BAG3 contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner (provided by RefSeq, Jul 2008).
	Goat anti Human BAG3 antibody detects a band of 85 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti BAG3 detects a band of approximately 85 kDa in MCF7 cell lysates.
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg MCF7 lysate lyophilized in RIPA buffer.
Lysate Reconstitution	<ul> <li>If using DDT reconstitute the lyophilized lysate with 190μl DI H<sub>2</sub>O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT.</li> <li>If using BME reconstitute the lyophilized lysate with 180μl DI H<sub>2</sub>O, add 200μl 2x Laemmli</li> </ul>
	Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf</a> Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a>

# **Related Products**

# **Recommended Secondary Antibodies**

Donkey Anti Sheep IgG (STAR88...) HRP

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

**America** Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

 Email: antibody\_sales\_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M351836:190318'

#### Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint