

Datasheet: VPA00123KT

Description:	CCT ALPHA ANTIBODY WITH CONTROL LYSATE
Specificity:	CCT ALPHA
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Specificity	Goat anti Human CCT alpha antibody recognizes the CCT alpha protein, also known as		
Synonyms	CCT1, CCTA		
	Entrez Gene: 6950 TCP1 Related reagents		
External Database Links	UniProt: P17987 Related reagents		
Immunogen	Peptide with the sequence C-SYEDAVHSGALND, from the C Terminus of the protein sequence.		
Preservative Stabilisers	0.02% Sodium Azide (NaN ₃) 0.5 % BSA		
Buffer Solution	TRIS buffered saline		
Preparation	20µl Goat polyclonal antibody purified by affinity chromatography		
Product Form	Purified IgG - liquid		
Target Species	Human		

T-complex protein 1 subunit alpha (TCP-1-alpha).

CCT alpha, encoded by the TCP1 gene, is a molecular chaperone that is a member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin. Alternate transcriptional splice variants of TCP1, encoding different isoforms, have been characterized. In addition, three pseudogenes that appear to be derived from TCP1 have been found (provided by RefSeq, Jun 2010).

Goat anti Human CCT alpha antibody detects a band of 60 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti CCT alpha detects a band of approximately 60 kDa in K562 cell lysates.	
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.	
Lysate Composition	400μg K562 lysate lyophilized in RIPA buffer.	
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H ₂ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT.	
	- If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME.	
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.	
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.	
Guarantee	As supplied, 12 months from date of despatch.	
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.	
Health And Safety Information	Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	
Regulatory	For research purposes only	

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

51

Worldwide

Tel: +44 (0)1865 852 700 **Europe**

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M348792:190228'

Printed on 15 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint