Datasheet: VPA00125 BATCH NUMBER 161214

Description:	GOAT ANTI HEXIM1
Specificity:	HEXIM1
Format:	Purified
Product Type:	PrecisionAb Polyclonal
lsotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
	rad-antibodies.com/prot	Yes	No	Not Determined	Suggested Dilution			
	Western Blotting	•			1/1000			
	criteria within Bio-Rac how we validate our P	I's ongoin recisionA nique this o	g antibo . b range . does not r	dy validation progran Where this product ha necessarily exclude its	e defined performance nme. Click <u>here</u> to learn s not been tested for use in such procedures.			
Target Species	Human							
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.							
Product Form	Purified IgG - liquid							
Preparation	Goat polyclonal antibod	y purified	by affinity	chromatography.				
Buffer Solution	TRIS buffered saline							
Preservative Stabilisers	0.02% Sodium Azide (N	laN ₃) 0.5 9	% BSA					

Immunogen	Peptide with the sequence C-HRQQERAPLSKFGD, from the C Terminus of the protein sequence.			
External Database Links	UniProt: 094992 Related reagents			
	Entrez Gene:			
	10614 HEXIM1 Related reagents			
Synonyms	CLP1, EDG1, HIS1, MAQ1			
Specificity	Goat anti Human HEXIM1 antibody recognizes protein HEXIM1, also known as cardiac lineage protein 1, estrogen down-regulated gene 1 protein, hexamethylene bis-acetamide-inducible protein 1 or menage a quatre protein 1.			
	Expression of HEXIM1 is induced by hexamethylene-bis-acetamide in vascular smooth muscle cells. HEXIM1 has no introns (provided by RefSeq, Jul 2008).			
	Goat anti Human HEXIM1 antibody detects a band of 70 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.			
Western Blotting	Anti HEXIM1 detects a band of approximately 70 kDa in K562 cell lysates.			
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.			
Guarantee	12 months from date of despatch			
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.			
Health And Safety Information	Material Safety Datasheet documentation #10058 available at: https://www.bio-rad-antibodies.com/SDS/VPA00125 Antibody (10058)			
Regulatory	For research purposes only			

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad	.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M370515:200529'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint