

Datasheet: VPA00127KT

Description:	ALOX15 ANTIBODY WITH CONTROL LYSATE		
Specificity:	ALOX15		
Format:	Purified		
Product Type:	PrecisionAb Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	2 Westerns		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
	Western Blotting	•			1/1000	
	PrecisionAb antibodies application. The antibod product has not been tes exclude its use in such p sample type.	ly has bee sted for us	en validato e in a par	ed at the suggested dilution the suggested di	ution. Where this oes not necessarily	
Target Species	Human					
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG - liquid					
Preparation	20µl Goat polyclonal antibody purified by affinity chromatography					
Buffer Solution	TRIS buffered saline					
Preservative Stabilisers	0.02% Sodium Azide (NaN ₃) 0.5% BSA					
Immunogen	Peptide with the sequence	ce C-HYK	TDVAVK	DDPE, from the internal	l region of the protein	

External Databases UniProt: P16050 Related reagents Einks P16050 Related reagents Synonyms LOG15 Specificity Goat anti Human ALOX15 antibody recognizes the Arachidonate 15-lipoxygenase, also known as 15-LOX, 15-LOX-1, 12/15-lipoxygenase, leukocyte-type Arachidonate 12-lipoxygenase, 12-LOX or Arachidonate omega-6 lipoxygenase. Goat anti Human ALOX15 antibody detects a band of 70 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Reconstitution Youg HEK293 lysate lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmil Sample Buffer and 10µl 2M DTT. Lysate Reconstitution Store lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmil Sample Buffer and 10µl 2M DTT. Syning BME reconstitute the lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmil Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C. Are reconstitution aliquot and store at -20°C for up to yain yain yain yain yain yain yain yain		sequence		
Specificity Goat anti Human ALOX15 antibody recognizes the Arachidonate 15-lipoxygenase, also known as 15-LOX, 15-LOX-1, 12/15-lipoxygenase, leukocyte-type Arachidonate 12-lipoxygenase, 12-LOX or Arachidonate omega-6 lipoxygenase. Goat anti Human ALOX15 antibody detects a band of 70 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Anti ALOX15 detects a band of approximately 70 kDa in HEK293 cell lysates. Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer. Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories.		P16050 Related reagents Entrez Gene:		
known as 15-LOX, 15-LOX-1, 12/15-lipoxygenase, leukocyte-type Arachidonate 12-lipoxygenase, 12-LOX or Arachidonate omega-8 lipoxygenase. Goat anti Human ALOX15 antibody detects a band of 70 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Anti ALOX15 detects a band of approximately 70 kDa in HEK293 cell lysates. Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer. Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb [™] is a trademark of Bio-Rad Laboratories. Heatth And Safety Information (10058): https://www.bio-rad-antibodies	Synonyms	LOG15		
Western Blotting Anti ALOX15 detects a band of approximately 70 kDa in HEK293 cell lysates. Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer. Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Specificity	known as 15-LOX , 15-LOX-1, 12/15-lipoxygenase, leukocyte-type Arachidonate 12-lipoxygenase, 12-LOX or Arachidonate omega-6 lipoxygenase. Goat anti Human ALOX15 antibody detects a band of 70 kDa. The antibody has been		
Instructions For Use Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer. Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf		extensively validated for western blotting using whole cell lysates.		
secondary antibody dilution and exposure time see product web page. Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer. Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf	Western Blotting	Anti ALOX15 detects a band of approximately 70 kDa in HEK293 cell lysates.		
Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Instructions For Use			
Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf	Lysate Composition	400µg HEK293 lysate lyophilized in RIPA buffer.		
Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Lysate Reconstitution	Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb		
up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch. Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		
Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>				
Health And Safety Material Safety Datasheet documentation #10058 #10561 available at: Information Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf	Guarantee	As supplied, 12 months from date of despatch.		
Information Antibody (10058): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>	Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.		
Regulatory For research purposes only		Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf		
	Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bic	-rad.com	Email: antibody_sales_de@bio-rad.com
		45 0004	and the second		

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351895:190318'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint