

Datasheet: VPA00130KT

Description:	POU2F1 ANTIBODY WITH CONTROL LYSATE
Specificity:	POU2F1
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Specificity	Goat anti Human POU2F1 antibody recognizes the POU domain, class 2, transcription factor 1,
Synonyms	OCT1, OTF1
	Entrez Gene: <u>5451</u> POU2F1 Related reagents
External Database Links	UniProt: P14859 Related reagents
Immunogen	Peptide with the sequence C-NVQSKSNEESGDSQQ, from the internal region of the protein sequence
Preservative Stabilisers	0.02% Sodium Azide (NaN ₃) 0.5% BSA
Buffer Solution	TRIS buffered saline
Preparation	20μl Goat polyclonal antibody purified by affinity chromatography
Product Form	Purified IgG - liquid
Target Species	Human

also known as NF-A1 Octamer-binding protein 1, Oct-1 Octamer-binding transcription factor 1 or

OTF-1.

The OCT1 transcription factor was among the first identified members of the POU transcription factor family (summarized by Sturm *et al.* 1993 (PubMed 8314572)). Members of this family contain the POU domain, a 160-amino acid region necessary for DNA binding to the octameric sequence ATGCAAAT (supplied by OMIM, Jul 2010).

Goat anti Human POU2F1 antibody detects a band of 94 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Anti POU2F1 detects a band of approximately 94 kDa in Ramos cell lysates.
Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
400μg Ramos lysate lyophilized in RIPA buffer.
- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H ₂ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT.
- If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME.
Heat at 95°C for 5 minutes. For 10 well mini gels load 25 μ l. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
As supplied, 12 months from date of despatch.
PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf

Related Products

Regulatory

America

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) HRP

North & South Tel: +1 800 265 7376

Fax: +1 919 878 3751

Worldwide

For research purposes only

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

'M348889:190228'

Fax: +44 (0)1865 852 739

Printed on 15 Mar 2019