

## Datasheet: VPA00131KT

<b>Description:</b>	CRTC2 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	CRTC2
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb Polyclonal
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	2 Westerns

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	20µl Goat polyclonal antibody purified by affinity chromatography
<b>Buffer Solution</b>	TRIS buffered saline
<b>Preservative Stabilisers</b>	0.02% Sodium Azide (NaN <sub>3</sub> ) 0.5 % BSA
<b>Immunogen</b>	Peptide with the sequence C-DPAVEDSFRSDRLQ, from the C Terminus of the protein

sequence

---

**External Database****Links****UniProt:**[Q53ET0](#)[Related reagents](#)**Entrez Gene:**[200186](#)

CRTC2

[Related reagents](#)

---

**Synonyms**

TORC2

---

**Specificity**

**Goat anti Human CRTC2 antibody** recognizes the CREB-regulated transcription coactivator 2, also known as transducer of regulated cAMP response element-binding protein 2 (TORC-2) or transducer of CREB protein 2.

CRTC2 is a member of the transducers of regulated cAMP response element-binding protein activity family of transcription coactivators. These proteins promote the transcription of genes targeted by the cAMP response element-binding protein, and therefore play an important role in many cellular processes. Under basal conditions CRTC2 is phosphorylated by AMP-activated protein kinase or the salt-inducible kinases and is sequestered in the cytoplasm. Upon activation by elevated cAMP or calcium, CRTC2 translocates to the nucleus and increases target gene expression. Single nucleotide polymorphisms in the CRTC2 gene may increase the risk of type 2 diabetes. A pseudogene of CRTC2 is located on the long arm of chromosome 5 (provided by RefSeq, Dec 2010).

Goat anti Human CRTC2 antibody detects a band of 80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

---

**Western Blotting**

Anti CRTC2 detects a band of approximately 80 kDa in MCF7 cell lysates.

---

**Instructions For Use**

Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

---

**Lysate Composition**

400µg MCF7 lysate lyophilized in RIPA buffer.

---

**Lysate Reconstitution**

- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.  
- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.  
Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

---

**Storage**

Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.

---

<b>Guarantee</b>	As supplied, 12 months from date of despatch.
<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf</a> Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) [HRP](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M351871:190318'

**Printed on 09 Feb 2021**