

Datasheet: VPA00155KT

Description:	PSMB8 ANTIBODY WITH CONTROL LYSATE
Specificity:	PSMB8
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species

Human

Species Cross Reactivity

Reacts with: Rat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

20µl Goat polyclonal antibody purified by affinity chromatography

Buffer Solution

TRIS buffered saline

Preservative Stabilisers

0.02% Sodium Azide (NaN₃) 0.5% BSA

Immunogen

Peptide with the sequence C-DVSDLLHQYREANQ, from the C terminus of the protein

sequence

External Database**Links****UniProt:**[P28062](#)[Related reagents](#)**Entrez Gene:**[5696](#)

PSMB8

[Related reagents](#)

Synonyms

LMP7, PSMB5i, RING10, Y2

Specificity

Goat anti Human PSMB8 antibody recognizes proteasome subunit beta type-8, also known as Low molecular mass protein 7, macropain subunit C13, multicatalytic endopeptidase complex subunit C13, proteasome component C13, and proteasome subunit beta-5i.

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMB8 encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. PSMB8 is located in the class II region of the MHC (major histocompatibility complex). Expression of PSMB8 is induced by gamma interferon and PSMB8 product replaces catalytic subunit 3 (proteasome beta 5 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to yield the same mature subunit (provided by RefSeq, Jul 2008).

Goat anti Human PSMB8 antibody detects a band of 23 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting

Anti PSMB8 detects a band of approximately 23 kDa in Jurkat cell lysates.

Instructions For Use

Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Lysate Composition

400µg Jurkat lysate lyophilized in RIPA buffer.

Lysate Reconstitution

- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.
- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.
Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M351911:190318'

Printed on 09 Feb 2021