

## Datasheet: VPA00171KT

<b>Description:</b>	HISTONE DEACETYLASE 6 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	HISTONE DEACETYLASE 6
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb Polyclonal
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	2 Westerns

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	20µl Rabbit polyclonal antibody purified by affinity chromatography
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin <30% Glycerol
<b>Immunogen</b>	Synthetic peptide corresponding to amino acids close to the N terminus of human histone deacetylase 6

#### External Database Links

##### UniProt:

[Q9UBN7](#)

[Related reagents](#)

**Entrez Gene:**

[10013](#) HDAC6 [Related reagents](#)

---

<b>Synonyms</b>	KIAA0901
-----------------	----------

---

<b>Specificity</b>	<p><b>Rabbit anti Human histone deacetylase 6 antibody</b> recognizes histone deacetylase 6, also known as HDAC6.</p> <p>Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Encoded by HDAC6 gene, histone deacetylase 6 belongs to class II of the histone deacetylase/acuc/apha family. It contains an internal duplication of two catalytic domains which appear to function independently of each other. This protein possesses histone deacetylase activity and represses transcription (provided by RefSeq, Jul 2008).</p> <p>Rabbit anti Human histone deacetylase 6 antibody detects a band of 160 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
--------------------	--

---

<b>Western Blotting</b>	Anti histone deacetylase 6 detects a band of approximately 160 kDa in HEK293 cell lysates.
-------------------------	--

---

<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
-----------------------------	---

---

<b>Lysate Composition</b>	400µg HEK293 lysate lyophilized in RIPA buffer.
---------------------------	---

---

<b>Lysate Reconstitution</b>	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
------------------------------	--

---

<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
----------------	--

---

<b>Guarantee</b>	As supplied, 12 months from date of despatch.
------------------	---

---

<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
-------------------------	--

---

<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf</a> Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a></p>
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](http://bio-rad-antibodies.com)

'M351169:190315'

**Printed on 09 Feb 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)