

Datasheet: VPA00196KT

| Description:  | GLI2 ANTIBODY WITH CONTROL LYSATE |
|---------------|-----------------------------------|
| Specificity:  | GLI2                              |
| Format:       | Purified                          |
| Product Type: | PrecisionAb Polyclonal            |
| Isotype:      | Polyclonal IgG                    |
| Quantity:     | 2 Westerns                        |
|               |                                   |

### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                  | Yes | No | Not Determined | Suggested Dilution |
|------------------|-----|----|----------------|--------------------|
| Western Blotting | •   |    |                | 1/1000             |

**PrecisionAb antibodies have been extensively** <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

| Buffer Solution Preservative | Phosphate buffered saline  0.09% Sodium Azide (NaN <sub>3</sub> )              |
|------------------------------|--|
| Stabilisers                  | 1% Bovine Serum Albumin  |
| Immunogen                    | Synthetic peptide corresponding to amino acids 46-60 of human GLI2 (isoform a) |
| External Database            |  |

Related reagents

**Entrez Gene:** 

P10070

| Synonyms              | THP   |
|-----------------------|---|
| Specificity           | Rabbit anti Human GLI2 antibody recognizes GLI2, also known as GLI family zinc finger protein 2, GLI-Kruppel family member GLI2, glioma-associated oncogene family zinc finger 2, oncogene GLI2, tax helper protein 1, tax helper protein 2, tax-responsive element-2 holding protein, tax-responsive element-25-bp sequence binding protein and zinc finger protein GLI2.  |
|                       | The GLI2 gene encodes a protein which belongs to the C2H2-type zinc finger protein subclass of the Gli family. Members of this subclass are characterized as transcription factors which bind DNA through zinc finger motifs. These motifs contain conserved H-C links. Gli family zinc finger proteins are mediators of Sonic hedgehog (Shh) signaling and they are implicated as potent oncogenes in the embryonal carcinoma cell. The protein encoded by GLI2 localizes to the cytoplasm and activates patched Drosophila homolog (PTCH) gene expression. It is also thought to play a role during embryogenesis. The encoded protein is associated with several phenotypes- Greig cephalopolysyndactyly syndrome, Pallister-Hall syndrome, preaxial polydactyly type IV, postaxial polydactyly types A1 and B (provided by RefSeq, Jul 2008). |
|                       | Rabbit anti Human GLI2 antibody detects a band of 191 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.   |
| Western Blotting      | Anti GLI2 detects a band of approximately 191 kDa in HeLa cell lysates.   |
| Instructions For Use  | Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.   |
| Lysate Composition    | 400μg HeLa lysate lyophilized in RIPA buffer.   |
| Lysate Reconstitution | - If using DDT reconstitute the lyophilized lysate with 190μl DI H <sub>2</sub> O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT If using BME reconstitute the lyophilized lysate with 180μl DI H <sub>2</sub> O, add 200μl 2x Laemmli Sample Buffer and 20μl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.  |
| Storage               | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.   |
|                       | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.  |
| Guarantee             | As supplied, 12 months from date of despatch.   |
| Acknowledgements      | PrecisionAb™ is a trademark of Bio-Rad Laboratories.  |
| Health And Safety     | Material Safety Datasheet documentation #10041 #10561 available at:   |

Information Antibody (10041): https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf

Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf

Regulatory For research purposes only

# **Related Products**

# **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376 America

Worldwide Tel: +44 (0)1865 852 700 Europe

Tel: +49 (0) 89 8090 95 21

Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351108:190315'

#### Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint