

Datasheet: VPA00217

Description:	RABBIT ANTI TISSUE TRANSGLUTAMINASE
Specificity:	TISSUE TRANSGLUTAMINASE
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Rabbit Ig fraction prepared by ammonium sulphate precipitation.
Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃).
Immunogen	KLH conjugated synthetic peptide between 429-458 amino acids from the Central region of human tissue transglutaminase

External Database Links

UniProt:

[P21980](#)

[Related reagents](#)

Entrez Gene:

Specificity **Rabbit anti Human tissue transglutaminase antibody** recognizes tissue transglutaminase, also known as protein-glutamine gamma-glutamyltransferase 2, C polypeptide, TG(C), TGase C, TGase H, TGase-2, TGase-H, epididymis secretory protein Li 45, transglutaminase C, transglutaminase H and transglutaminase-2.

Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. Encoded by the TGM2 gene, tissue transglutaminase acts as a monomer, is induced by retinoic acid, and appears to be involved in apoptosis. It is also the autoantigen implicated in celiac disease. Two transcript variants encoding different isoforms have been found for TGM2 (provided by RefSeq, Jul 2008).

Rabbit anti Human tissue transglutaminase antibody detects a band of 78 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Anti TGM2 detects a band of approximately 78 kDa in K562 cell lysates.

Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Guarantee 12 months from date of despatch.

Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/VPA00217>
Antibody (10040)

Regulatory For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M400850:220712'

Printed on 19 Jan 2024