

Datasheet: VPA00269

BATCH NUMBER 160415

Description:	RABBIT ANTI FGR
Specificity:	FGR
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Ig fraction - liquid
Preparation	Rabbit Ig fraction prepared by ammonium sulphate precipitation
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	KLH-conjugated synthetic peptide corresponding to aa 3-33 of human FGR
External Database Links	UniProt: P09769 Related reagents

Entrez Gene:[2268](#) FGR [Related reagents](#)

Synonyms	SRC2
-----------------	------

Specificity	<p>Rabbit anti Human FGR antibody recognizes FGR, also known as c-src-2 proto-oncogene, p55-c-fgr protein, proto-oncogene c-Fgr, tyrosine-protein kinase Fgr and v-fgr feline Gardner-Rasheed sarcoma viral oncogene homolog.</p> <p>FGR is a member of the Src family of protein tyrosine kinases (PTKs). The FGR gene encoded protein contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of FGR. Multiple alternatively spliced variants, encoding the same protein, have been identified (provided by RefSeq, Jul 2008).</p> <p>Rabbit anti Human FGR antibody detects a band of 56 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
--------------------	---

Western Blotting	Anti FGR detects a band of approximately 56 kDa in Ramos cell lysate
-------------------------	--

Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
-------------------------	---

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VPA00269 Antibody (10040)
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M370631:200529'

Printed on 25 Mar 2023

