

Datasheet: VPA00269KT

Description:	FGR ANTIBODY WITH CONTROL LYSATE		
Specificity:	FGR		
Format:	Purified		
Product Type:	PrecisionAb™ Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	2 Westerns		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

PrecisionAb antibodies have been extensively validated for the western blot application. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human		
Product Form	lg fraction - liquid		
Preparation	20μl Rabbit Ig fraction prepared by ammonium sulphate precipitation		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)		
Immunogen	KLH-conjugated synthetic peptide corresponding to aa 3-33 of human FGR		
External Database Links	UniProt: P09769 Related reagents Entrez Gene: 2268 FGR Related reagents		
Synonyms	SRC2		
Specificity	Rabbit anti Human FGR antibody recognizes FGR, also known as c-src-2		

c-2 proto-oncogene, p55-c-fgr protein, proto-oncogene c-Fgr, tyrosine-protein kinase Fgr and v-fgr feline Gardner-

Rasheed sarcoma viral oncogene homolog.

FGR is a member of the Src family of protein tyrosine kinases (PTKs). The FGR gene encoded protein contains N-terminal sites for myristylation and palmitylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosinecontaining and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of FGR. Multiple alternatively spliced variants, encoding the same protein, have been identified (provided by RefSeq, Jul 2008).

Rabbit anti Human FGR antibody detects a band of 56 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti FGR detects a band of approximately 56 kDa in Ramos cell lysate
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg Ramos lysate lyophilized in RIPA buffer
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT.
	- If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME.
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Shelf Life	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety	Material Safety Datasheet documentation #10040 #10561 available at:
Information	Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
	Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Worldwide

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M337021:181214'

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint