

Datasheet: VPA00280

BATCH NUMBER 160415

Description:	RABBIT ANTI LIMK2
Specificity:	LIMK2
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 μΙ

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	-			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human	
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary betwee reactivity is derived from testing within our laboratories, peer-repersonal communications from the originators. Please refer to further information.	eviewed publications or
Product Form	Purified IgG - liquid	
Preparation	Rabbit polyclonal antibody purified by affinity chromatography	
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)	

Immunogen

KLH-conjugated synthetic peptide corresponding to aa 570-599 of human LIMK2

External Database Links

UniProt:

P53671 Related reagents

Entrez Gene:

3985 LIMK2 Related reagents

Specificity

Rabbit anti Human LIMK2 antibody recognizes LIMK2, also known as LIM domain kinase 2.

There are approximately 40 known eukaryotic LIM proteins, named after the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by the LIMK2 gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for LIMK2 (provided by RefSeq, Jul 2008).

Rabbit anti Human LIMK2 antibody detects a band of 70 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti LIMK2 detects a band of approximately 70 kDa in HEK293 cell lysate
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles
Guarantee	12 months from date of despatch
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VPA00280 Antibody (10040)
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Europe

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint