

Datasheet: VPA00406KT

Description:	FIP1L1 ANTIBODY WITH CONTROL LYSATE
Specificity:	FIP1L1
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Mouse</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	20µl Rabbit polyclonal antibody purified by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<p>0.09% Sodium Azide (NaN₃)</p> <p>2% Sucrose</p>
Immunogen	Synthetic peptide corresponding to the C-terminal region of human FIP1L1

External Database Links	UniProt: Q6UN15 Related reagents
	Entrez Gene: 81608 FIP1L1 Related reagents
Synonyms	FIP1, RHE
Specificity	<p>Rabbit anti Human FIP1L1 antibody recognizes FIP1L1, also known as Pre-mRNA 3'-end-processing factor FIP1, FIP1-like 1 protein and factor interacting with PAP.</p> <p>The FIP1L1 gene encodes a subunit of the CPSF (cleavage and polyadenylation specificity factor) complex that polyadenylates the 3' end of mRNA precursors. FIP1L1, the homolog of yeast Fip1 (factor interacting with PAP), binds to U-rich sequences of pre-mRNA and stimulates poly(A) polymerase activity. Its N-terminus contains a PAP-binding site and its C-terminus an RNA-binding domain. An interstitial chromosomal deletion on 4q12 creates an in-frame fusion of human genes FIP1L1 and PDGFRA (platelet-derived growth factor receptor, alpha). The FIP1L1-PDGFRA fusion gene encodes a constitutively activated tyrosine kinase that joins the first 233 amino acids of FIP1L1 to the last 523 amino acids of PDGFRA. FIP1L1 fusion and chromosomal deletion is the cause of some forms of idiopathic hypereosinophilic syndrome (HES). This syndrome, recently reclassified as chronic eosinophilic leukemia (CEL), is responsive to treatment with tyrosine kinase inhibitors. Alternative splicing results in multiple transcript variants encoding distinct isoforms (provided by RefSeq, Oct 2008).</p> <p>Rabbit anti Human FIP1L1 antibody detects a band of 77 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
Western Blotting	Anti FIP1L1 detects a band of approximately 77 kDa in Jurkat cell lysate
Instructions For Use	Please refer to the PrecisionAb western blotting protocol . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg Jurkat lysate lyophilized in RIPA buffer
Lysate Reconstitution	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
Storage	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>

Guarantee As supplied, 12 months from date of despatch

Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories

Health And Safety Information Material Safety Datasheet documentation #10045 #10561 available at:
Antibody (10045): <https://www.bio-rad-antibodies.com/uploads/MSDS/10045.pdf>
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M351092:190315'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)