

Datasheet: VPA00407KT

Description:	TIMM50 ANTIBODY WITH CONTROL LYSATE			
Specificity:	TIMM50			
Format:	Purified			
Product Type:	PrecisionAb Polyclonal			
lsotype:	Polyclonal IgG			
Quantity:	2 Westerns			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>							
	rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution							
	Western Blotting	•	110	Not Determined	1/1000			
	PrecisionAb antibodie application. The antibo product has not been te exclude its use in such p sample type.	dy has bee sted for us	en validat e in a par	ed at the suggested dil ticular technique this d	ution. Where this loes not necessarily			
Target Species	Human							
Product Form	Purified IgG - liquid							
Preparation	20µl Rabbit polyclonal antibody purified by affinity chromatography							
Buffer Solution	Phosphate buffered saline							
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)							
Immunogen	KLH-conjugated synthet	tic peptide	correspoi	nding to aa 25-56 of hu	ıman TIMM50			
External Database Links	UniProt: Q3ZCQ8 Related	d reagents						

Entrez Gene:

Synonyms	TIM50
Specificity	Rabbit anti Human TIMM50 antibody recognizes TIMM50, also known as Tim50-like protein, homolog of yeast Tim50 and mitochondrial import inner membrane translocase subunit TIM50.
	Rabbit anti Human TIMM50 antibody detects a band of 40 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti TIMM50 detects a band of approximately 40 kDa in HEK293 cell lysate
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg HEK293 lysate lyophilized in RIPA buffer
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for
	up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>
Regulatory	For research purposes only

92609 TIMM50 Related reagents

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint