

## Datasheet: VPA00415KT

<b>Description:</b>	HISTONE H4 (DIMETHYL K20) ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	HISTONE H4 (DIMETHYL K20)
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb Polyclonal
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	2 Westerns

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Mouse, Rat

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG - liquid

### Preparation

20µl Rabbit polyclonal antibody purified by affinity chromatography

### Buffer Solution

Phosphate buffered saline

### Preservative Stabilisers

0.09% Sodium Azide (NaN<sub>3</sub>)

### Immunogen

KLH-conjugated synthetic peptide corresponding to aa 1-30 of human histone H4

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P62805</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">554313</a> HIST2H4B <a href="#">Related reagents</a>
<b>Synonyms</b>	H4/A, H4/B, H4/C, H4/D, H4/E, H4/G, H4/H, H4/I, H4/J, H4/K, H4/M, H4/N, H4/O, H4F2, H4FA, H4FB, H4FC, H4FD, H4FE, H4FG, H4FH, H4FI, H4FJ, H4FK, H4FM, H4FN, H4FO, HIST2H4
<b>Specificity</b>	<p><b>Rabbit anti Human histone H4 (dimethyl K20) antibody</b> recognizes histone H4 (dimethyl K20).</p> <p>Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.</p> <p>Rabbit anti Human histone H4 (dimethyl K20) antibody detects a band of 15 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
<b>Western Blotting</b>	Anti histone H4 (dimethyl K20) detects a band of approximately 15 kDa in NIH/3T3 cell lysate
<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
<b>Lysate Composition</b>	400µg NIH/3T3 lysate lyophilized in RIPA buffer
<b>Lysate Reconstitution</b>	<ul style="list-style-type: none"> <li>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</li> <li>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</li> </ul> Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>

**Guarantee** As supplied, 12 months from date of despatch

---

**Acknowledgements** PrecisionAb™ is a trademark of Bio-Rad Laboratories

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 #10561 available at:  
Antibody (10040): <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>  
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M351188:190315'

**Printed on 10 Feb 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)