

Datasheet: VPA00511

Description:	GOAT ANTI CORTACTIN
Specificity:	CORTACTIN
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 μΙ

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	-			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human	
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications o personal communications from the originators. Please refer to references indicated for further information.	
Product Form	Purified IgG - liquid.	
Preparation	Goat polyclonal antibody purified by affinity chromatography.	
Buffer Solution	TRIS buffered saline.	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 0.5% Bovine Serum Albumin.	
Immunogen	Peptide sequence WKASAGHAVSIA-C	

External Database Links

UniProt:

Q14247 Related reagents

Entrez Gene:

2017 CTTN Related reagents

Synonyms

EMS1

Specificity

Goat anti Human cortactin antibody recognizes the Src substrate cortactin, also known as amplaxin or oncogene EMS1.

The CTTN gene is overexpressed in breast cancer and squamous cell carcinomas of the head and neck (Patel et al. 1996). The encoded protein is localized in the cytoplasm and in areas of the cell-substratum contacts. CTTN has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of CTTN contributes to tumor cell invasion and metastasis (Chuma et al. 2004). Three splice variants that encode different isoforms have been identified for CTTN (provided by RefSeq, May 2010).

Goat anti Human cortactin antibody detects a band of 80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Anti cortactin detects a band of approximately 80 kDa in NIH/3T3 cell lysates.

Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Guarantee 12 months from date of despatch.

Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories.

Health And Safety Information

Material Safety Datasheet documentation #10058 available at:

https://www.bio-rad-antibodies.com/SDS/VPA00511
Antibody (10058)

Regulatory

For research purposes only.

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) HRP

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700 **Europe**

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421518:230706'

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint