

Datasheet: VPA00543KT

Description:	WILMS TUMOR 1 ANTIBODY WITH CONTROL LYSATE		
Specificity:	WILMS TUMOR 1		
Format:	Purified		
Product Type:	PrecisionAb™ Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	2 Westerns		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	-			1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human			
Product Form	Purified IgG - liquid.			
Preparation	20μl Rabbit polyclonal antibody purified by affinity chromatography.			
Buffer Solution	Phosphate buffered saline.			
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃).			
Immunogen	KLH conjugated synthetic peptide corresponding to amino acid 346-375 of human Wilm's tumor 1.			
External Database Links	UniProt: P19544 Related reagents			

Related reagents

Specificity

Entrez Gene:

WT1

Rabbit anti Human Wilm's tumor 1 antibody recognizes Wilms tumor 1, also known as WT1, amino-terminal domain of EWS or WT33.

WT1 gene encodes a transcription factor that contains four zinc-finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with

Wilm's tumors. WT1 exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation site upstream of and in-frame with the first AUG. WT1 mRNA undergoes RNA editing in human and rat, and this process is tissue-restricted and developmentally regulated (Sharma et al. 1994) (provided by RefSeq, Oct 2010).

Rabbit anti Human Wilm's tumor 1 antibody detects a band of 55 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Anti Wilms tumor 1 detects a band of approximately 55 kDa in K562	cell lysates.
Instructions For Use	Please refer to the PrecisionAb western blotting protocol.	
Lysate Composition	400μg K562 lysate lyophilized in RIPA buffer.	
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H ₂ O, Buffer and 10 μ l 2M DTT.	add 200µl 2x Laemmli Sample
	- If using BME reconstitute the lyophilized lysate with 180 μ l DI H ₂ O, Buffer and 20 μ l BME.	add 200µl 2x Laemmli Sample
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other	er gel and comb formats please
	refer to the PrecisionAb western blotting protocol.	
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cy	ycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot months or at -80°C for longer term storage.	and store at -20°C for up to 3
Guarantee	As supplied, 12 months from date of despatch.	
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.	
Health And Safety	Material Safety Datasheet documentation #10040 #10561 available	at:
Information	Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS	<u>8/10040.pdf</u>
	Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads	/MSDS/10561.pdf
Regulatory	For research purposes only.	

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376 Worldwide America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M339020:181217'

Printed on 16 Mar 2019