

Datasheet: VPA00573KT

Description:	PAXILLIN ANTIBODY WITH CONTROL LYSATE		
Specificity:	PAXILLIN		
Format:	Purified		
Product Type:	PrecisionAb Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	2 Westerns		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Rabbit polyclonal antibody purified by affinity chromatography.
Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃).
Immunogen	KLH conjugated synthetic peptide corresponding to 94-125 amino acid from human paxillin
External Database Links	UniProt:

Related reagents

P49023

Entrez Gene:

5829 PXN Related reagents

Specificity	Rabbit anti Human paxillin antibody recognizes paxillin.
	The PXN gene encodes a cytoskeletal protein involved in actin-membrane attachment at sites of cell adhesion to the extracellular matrix (focal adhesion). Alternatively spliced transcript variants encoding different isoforms have been described for PXN. These isoforms exhibit different expression pattern, and have different biochemical, as well as physiological properties (Mazaki et al. 1997).
	Rabbit anti Human paxillin antibody detects a band of 62 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti paxillin detects a band of approximately 62 kDa in A431 cell lysates.
Instructions For Use	Please refer to the PrecisionAb western blotting protocol.
Lysate Composition	400μg A431 lysate lyophilized in RIPA buffer.
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190μl DI H ₂ O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT If using BME reconstitute the lyophilized lysate with 180μl DI H ₂ O, add 200μl 2x Laemmli Sample Buffer and 20μl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21
 From

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50
 March

 Email: antibody_sales_us@bio-rad.com
 Email: antibody_sales_uk@bio-rad.com
 Email: antibody_sales_de@bio-rad.com
 Email: antibody_sales_de@bio-rad.com

2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351310:190318'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint