

Datasheet: VPA00587

Description:	RABBIT ANTI MEK 2
Specificity:	MEK 2
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 μΙ

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.biorad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

2

External Database	UniProt:
Immunogen	Synthetic peptide directed towards the N terminal region of human MEK
Stabilisers	2% Sucrose.
Preservative	0.09% Sodium Azide (NaN ₃)
Buffer Solution	Phosphate buffered saline.
Preparation	Rabbit polyclonal antibody purified by affinity chromatography.
Product Form	Purified IgG - liquid.
Target Species	Human

P36507 Related reagents

Entrez Gene:

5605 MAP2K2 Related reagents

Synonyms	MEK2, MKK2, PRKMK2
Specificity	Rabbit anti Human MEK 2 antibody recognizes MEK 2 also known as dual specificity mitogen-activated protein kinase kinase 2, also known as ERK activator kinase 2, MAP kinase kinase 2, MAPK/ERK kinase 2, MAPKK2 or p45.
	The protein encoded by MAP2K2 gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in MAP2K2 gene cause Cardiofaciocutaneous syndrome 4 (CFC4), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome (Schulz et al. 2008). The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for MAP2K2 (provided by RefSeq, Jul 2008).
Western Blotting	extensively validated for western blotting using whole cell lysates. Anti MEK 2 detects a band of approximately 49 kDa in HepG2 cell lysates.
-	
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
Guarantee	12 months from date of despatch.
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10045 available at: https://www.bio-rad-antibodies.com/SDS/VPA00587 Antibody (10045)
Regulatory	For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376
America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M402747:220719'

Printed on 13 Aug 2023