

Datasheet: VPA00631KT

Description:	p43 ANTIBODY WITH CONTROL LYSATE		
Specificity:	p43		
Format:	Purified		
Product Type:	PrecisionAb™ Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	2 Westerns		

Product Details

Applications

Specificity

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human		
Product Form	Purified IgG - liquid		
Preparation	20μl Rabbit polyclonal antibody purified by affinity chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
Immunogen	KLH conjugated synthetic peptide of partial human p43 (amino acids 342 - 370)		
External Database Links	UniProt: P49411 Related reagents Entrez Gene: 7284 TUFM Related reagents		

Rabbit anti Human p43 antibody recognizes the elongation factor Tu, mitochondrial, also known as EF-Tu or TUFM.

The TUFM gene encodes a protein which participates in protein translation in mitochondria. Mutations in TUFM have been associated with combined oxidative phosphorylation deficiency

Rabbit anti Human p43 antibody detects a band of 46 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti p43 detects a band of approximately 46 kDa in HepG2 cell lysates **Instructions For Use** Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page. **Lysate Composition** 400µg HepG2 lysate lyophilized in RIPA buffer. Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. Guarantee As supplied, 12 months from date of despatch **Acknowledgements** PrecisionAb™ is a trademark of Bio-Rad Laboratories **Health And Safety** Material Safety Datasheet documentation #10040 #10561 available at: Information Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf

resulting in lactic acidosis and fatal encephalopathy. A pseudogene has been identified on

chromosome 17 (provided by RefSeq, Jul 2008).

Related Products

Regulatory

Recommended Secondary Antibodies

Email: antibody sales us@bio-rad.com

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

For research purposes only

Tel: +44 (0)1865 852 700 **Europe**

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody sales de@bio-rad.com

'M337867:181217'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint